

# PATENT COOPERATION TREATY

PCT

## NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

United States Patent and Trademark  
Office  
(Box PCT)  
Crystal Plaza 2  
Washington, DC 20231  
ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

Date of mailing (day/month/year)

27 July 1998 (27.07.98)

International application No.

PCT/CA97/00969

Applicant's or agent's file reference

12771

International filing date (day/month/year)

15 December 1997 (15.12.97)

Priority date (day/month/year)

13 December 1996 (13.12.96)

Applicant

MUNROE, Donald, G. et al

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:

07 July 1998 (07.07.98)

☐ in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was

☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO  
34, chemin des Colombettes  
1211 Geneva 20, Switzerland

Facsimile No.: (41-22) 740.14.35

Authorized officer

S. Cruz

Telephone No.: (41-22) 338.83.38

## PATENT COOPERATION TREATY

PCT

REC'D 23 MAR 1999

WIPO PCT

## INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference 12771	<b>FOR FURTHER ACTION</b> See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/CA97/00969	International filing date (day/month/year) 15/12/1997	Priority date (day/month/year) 13/12/1996
International Patent Classification (IPC) or national classification and IPC C12N15/12		
Applicant ALLELIX BIOPHARMACEUTICALS INC. et al.		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.


2. This REPORT consists of a total of 5 sheets, including this cover sheet.

- ☒ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of 21 sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☒ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☐ Certain observations on the international application

Date of submission of the demand  07/07/1998	Date of completion of this report
Name and mailing address of the international preliminary examining authority:  European Patent Office D-80298 Munich Tel. (+49-89) 2399-0 Tx: 523656 epmu d Fax: (+49-89) 2399-4465	Authorized officer  van Heusden, M  Telephone No. (+49-89) 2399



# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/CA97/00969

## I. Basis of the report

1. This report has been drawn on the basis of (*substitute sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to the report since they do not contain amendments.*):

### Description, pages:

1,2,4-23,25-31,33, as originally filed  
35,37,39

3,24,32,34,36, as received on 20/04/1998 with letter of 17/04/1998  
38

### Claims, No.:

1-16 as received on 08/02/1999 with letter of 05/02/1999

### Drawings, sheets:

1/13-13/13 as received on 20/04/1998 with letter of 17/04/1998

2. The amendments have resulted in the cancellation of:

- ☐ the description, pages:  
☐ the claims, Nos.:  
☐ the drawings, sheets:

3. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

4. Additional observations, if necessary:

## II. Priority

1. ☐ This report has been established as if no priority had been claimed due to the failure to furnish within the prescribed time limit the requested:
- ☐ copy of the earlier application whose priority has been claimed.
- ☐ translation of the earlier application whose priority has been claimed.

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT**

International application No. PCT/CA97/00969

2. ☐ This report has been established as if no priority had been claimed due to the fact that the priority claim has been found invalid.

Thus for the purposes of this report, the international filing date indicated above is considered to be the relevant date.

3. Additional observations, if necessary:

**see separate sheet**

**V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement**

1. Statement

Novelty (N)	Yes:	Claims	1-16
	No:	Claims	
Inventive step (IS)	Yes:	Claims	
	No:	Claims	1-16
Industrial applicability (IA)	Yes:	Claims	1-16
	No:	Claims	

2. Citations and explanations

**see separate sheet**



**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/CA97/00969

**Additional remarks to section II:**

1. The priority document pertaining to the present application was not available at the time of establishing this IPER. Hence, the current assessment is based on the assumption that all claims enjoy priority rights from the filing date of the earliest priority document.

**Additional remarks to section V:**

**1. Citations**

- 1.1 The documents mentioned in this IPER are numbered as in the International Search Report (ISR), i.e. D1 corresponds to the first document of the ISR etc.

**2. Novelty (Article 33(2) PCT)**

- 2.1 The present application relates to an isolated polynucleotide encoding glucagon-like-protein 2 (GLP-2) receptor from human or from rat, to polynucleotides hybridizing to said receptor-encoding polynucleotides, and to the recombinant expression of said GLP-2 receptors. It further relates to antibodies to said receptors and to a method for identifying GLP-2 receptor ligands.
- 2.2 The present application satisfies the criterion set forth in Article 33(2) PCT in that the subject matter of claims 1-16 is novel with regard to the cited prior art.

**3. Inventive step (Article 33(3) PCT)**

- 3.1 The present application does not satisfy the criterion set forth in Article 33(3) PCT because the subject matter of claims 1-16 does not involve an inventive step over the disclosure of D1 in combination with the teaching e.g. of D3 for the following

## PATENT COOPERATION TREATY

USA

From the INTERNATIONAL SEARCHING AUTHORITY

PCT

To:

OSLER HOSKIN & HARCOURT  
Attn. AITKEN, D.W.  
50 O'Connor Street  
Suite 1500  
Ottawa Ontario K1P 6L2  
CANADA

COMMUNICATION IN CASES FOR WHICH  
NO OTHER FORM IS APPLICABLE

Date of mailing (day/month/year) 13/10/1998	
Applicant's or agent's file reference 12771	REPLY DUE See paragraph 1 below
International application No. PCT/CA 97/ 00969	International filing date (day/month/year) 15/12/1997
Applicant ALLELIX BIOPHARMACEUTICALS INC. et al.	

1. ☐ REPLY DUE within \_\_\_\_\_ ~~XXXX~~ days from the above date of mailing

☒ NO REPLY DUE

## 2. COMMUNICATION:

With reference to your letter of September 21st 1998, and concerning the International Search Report sent to you on June 10th 1998, where you inform us that Cited Document W098/25955 was not received by you in full.

It appears that in error, only a copy of the front page has been transmitted.

Please find enclosed a full copy of Cited Document W098/25955.

We apologize for any inconvenience caused.

Name and mailing address of the International Searching Authority



European Patent Office, P.B. 5818 Patentlaan 2  
NL-2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Andria Overbeeke-Siepkens

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/CA 97/00969

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C07K14/705 C12N5/10 C12N15/62 C07K16/28  
C12Q1/68 G01N33/50

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K C12N C12Q G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DRUCKER D J ET AL: "INDUCTION OF INTESTINAL EPITHELIAL PROLIFERATION BY GLUCAGON-LIKE PEPTIDE 2" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 93, 1990, pages 7911-7916, XP000650124	19
Y	whole document , especially page 7914, right column + page 7915	1-5,12, 13
X	--- WO 96 32414 A (1149336 ONTARIO INC) 17 October 1996 pages 2,3,6,7	19
Y	--- WO 93 19175 A (THORENS BERNARD) 30 September 1993 abstract, examples 1-3 ---	1-5,12, 13
-/-		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\* & \* document member of the same patent family

Date of the actual completion of the international search

12 October 1998

Date of mailing of the international search report

11 Oct. 98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Holtorf, S

## INTERNATIONAL SEARCH REPORT

Internat Application No

PCT/CA 97/00969

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 04821 A (MERCK & CO INC ;GRAZIANO MICHAEL P (US); BORKOWSKI DOREEN A (US);) 16 February 1995 pages 1-3, page 5 , line 27-33; page 6; example 5 -----	1-19

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/CA97/00969

reasons.

The closest prior art to evaluate the inventiveness of claims 1-16 is document D1, which discloses that the GLP-2 peptide stimulates intestinal epithelial proliferation. The authors of D1 (see sentence bridging p. 7914 to 7915) hypothesize that their 'observations may be explained by tissue specific expression of the **receptor for GLP-2**' and continue to suggest the isolation of the receptor for GLP-2. D1 thus suggests that **intestinal tissue** may express a GLP-2 receptor. Document D3 discloses the cloning of rat and human GLP-1 receptor, using standard techniques of recombinant DNA technology. Thus the prior art suggests both the existence and the location of a GLP-2 receptor (D1) and discloses a method on how to isolate the closely related GLP-1 receptor by recombinant DNA technology (D3), without isolating the actual receptor protein or knowing the amino acid sequence. Therefore it would be obvious for the skilled person to combine the teachings of D3 with the information disclosed in D1 in order to arrive at the cloning of the rat and/or human receptor for GLP-2, by applying standard hybridization and amplification techniques, which is therefore not considered to involve an inventive step.

**4. Industrial applicability (Article 33(4) PCT)**

The subject matter of claims 1-16 is considered to be industrially applicable.

Figure 4 discloses a CDNA sequence of 667 nucleotides (SEQ ID NO: 9) which encodes a 222 amino acid fragment (SEQ ID NO: 10) of a human GLP-2 receptor.

5 Figure 5 discloses the amino acid sequence (SEQ ID NO: 10) expressed from the CDNA of Figure 4.

Figure 6 discloses a cDNA sequence (SEQ ID NO: 11), nucleotides 121-1779 of which encode a human GLP-2 receptor.

10 Figure 7 discloses the amino acid sequence of the expression product (SEQ ID NO: 12) from the cDNA of Figure 6.

Figure 8 illustrates the functional activation by GLP-2 peptide of the human receptor encoded by SEQ ID NO: 11 (Figure 8).

15 Figure 9 compares the amino acid sequences of the rat GLP-2 receptor and the human GLP-2 receptor.

20 Figure 10 compares the amino acid sequences of the rat GLP-2 receptor and the human GLP-2 receptor against rat GLP-1 receptor (SEQ ID NO:13).

#### DETAILED DESCRIPTION AND PREFERRED EMBODIMENTS

25 The invention relates in one respect to polynucleotides, in their isolated form, that code for GLP-2 receptors. As used herein "isolated" means separated from polynucleotides that encode other proteins. In the context of polynucleotide libraries, for instance, GLP-2 receptor-encoding polynucleotide is considered "isolated" when it has been selected, and hence removed from association with other polynucleotides within the library. Such polynucleotides may be in the form of RNA, or in the form of DNA including cDNA, genomic DNA and synthetic DNA. The GLP-2 receptors are characterized by structural  
30 features common to the G-protein coupled receptor class, including seven transmembrane regions, and by the functional properties of binding GLP-2 peptide selectively, i.e.,

amplified from clonally impure plugs or 2777-clone pools. The sequences of the vector primers were as follows:

830F: [5'-AACCCACTGCTTAC-3'] (SEQ ID NO:14)

1186R: [5'-CCCAGAATAGAATGACACC-3'] (SEQ ID NO:15)

5 The PCR was done, under the following conditions just noted, using Expand™ PCR system.

10 The most prominent band was reamplified, purified and sequenced. Based on the amplified sequence obtained, additional primers were designed and new sequencing carried out. In this manner the complete sequences of the GLP-2R cDNA inserts in all five sources of clones were determined. Sequence analysis showed that only pool RHT 440 and pool RHT 587 contain clones with complete coding sequence of GLP-2R and that the two clones were identical (derived from the same cDNA clone).

15 Because of difficulty in clonally purifying the GLP-2 receptor cDNA clone from the RHT 440 or RHT 587 cDNA library pools, the cDNA was amplified and recloned into pcDNA3. Based on the sequence obtained from RUT 440 and RHT 587, two primers were designed one which primed starting 4bp upstream of the initiation codon and another which primed starting 8bp downstream of the stop codon.

20

WBR-C5: [5'-CAGGGGCCGGTACCTCTCCACTCC-3'] (SEQ ID NO:16)

WBR-C3: [5'-TTGGGTCCTCGAGTGGCCAAGCTGCG-3'] (SEQ ID NO:17)

25 The two primers were used to amplify a DNA fragment of approximately 1525 bp fragment under the following PCR conditions using Expand™ PCR system from Boehringer Mannheim (Catalogue no. 1681-842).

10 µl of 10x Expand™ PCR Buffer 1

14 µl of 2.5mM dNTP mix

3.0 µl of Primer 1 (10 µM) (WBR-C5)

30 3.0 µl of Primer 2 (10 µM) (WBR-C3)

1.5 µl of Enzyme (5 units)

the other is the universal amplification primer provided with the kit. The amplified product is subcloned into a plasmid vector for sequencing.

5 For the 5' RACE System, the first strand cDNA is synthesized from mRNA using a gene-specific primer (which is based on the available partial cDNA sequence of HHT13) and SuperScript II reverse transcriptase. The original mRNA template is removed by treatment with RNase H. Unincorporated dNTPs, primer, and proteins are separated from cDNA using spin cartridge. A homopolymeric dCTP tail is then added to the 3'- end of the first strand cDNA using TdT enzyme and dCTP nucleotides. PCR amplification is performed using two primers: one is a nested, gene-specific primer designed from the available partial DNA sequence of HHT13; and the other is an "anchor primer" provided with the system. Both primers incorporate restriction sites for subcloning into plasmids and subsequent sequencing.

15 Sub-cloning of HHT13  $\pi$ gt10 clones into pcdna3, their sequencing and expression

A. Amplification of cDNA inserts with  $\pi$ gt10 primers.

On a lawn of bacterial cells (E.coli C600Hf1), 10  $\mu$ l of phage resuspension from each clone was placed at marked spots. After 5 hr incubation at 37°C, the phage plaques were clearly visible. The surface of each plaque was transferred to 200  $\mu$ l of water. The samples were kept in boiling water bath for 5 minutes and centrifuged at room temperature for 10 minutes. 1  $\mu$ l of sample was used to amplify with a set of  $\pi$ gt10 primers.

20 GT10-5KXb [5'-GGGTAGTCGGTACCTCTAGAGCAAGTTCAGCC-3'] (SEQ ID NO:18)

25 vs

GT10-3BXh [5'-ATAACAGAGGATCCTCGAGTATTTCTTCCAG-3'] (SEQ ID NO:19)

The Expand™ PCR system from Boehringer Mannheim (Cat. 1681-842) was used under the following conditions:

30 5  $\mu$ l of 10x Expand™ Buffer 3

7  $\mu$ l of 2.5mM dNTP mix



D. Comparison of DNA sequences between rat GLP-2R and EEHT13 subclones. The comparison showed a 2 bp deletion at a position corresponding to nucleotides 389-390 of the rat GLP-2R cDNA, resulting in the loss of nucleotides 374-375 of the human GLP-2R cDNA sequence presented herein.

5

PCR was used to incorporate two bp of the rat GLP-2R DNA sequence into HHT13-1 DNA at the site of the 2 bp frame-shift deletion identified relative to the rat GLP-2R coding sequence. The following primers were designed from HHT13 DNA sequence to insert two bp

10

HWBR/2BPI-475F

[5'-ACAGGCATGTCTGGAAGACTTACTCAAGGAACCTTCTGGCAT-3']

(SEQ ID NO:20) HWBR/2BPI-506R

[5'-ATGCCAGAAGGTTCTTGTGAGTAAGTCTTCCAGACATGCCTGT-3'] (SEQ ID NO:21)

15

HWBR-F7 [5'-TTCCTCTGTGGTACCAAGAGGAATGC-3'] (SEQ ID NO:22)

and HWBR-1910R:

[5'-GGTGGACTCGAGGTACCGATCTCACTCTCTTCCAGAATC-3'] (SEQ ID NO:23)

20

PCR 1: One ng of pHHT13-1 DNA was used as template to do two PCRs with primers, HWBR-F7 vs HWBR/2BPI-506R and HWBR/2BPI-475F vs HWBR-1910R. The Expand™ PCR system from Boehringer Mannheim (Cat. 1681-842) was used under the following conditions:

5 µl of Lox Expand™ Buffer 3

7 µl of 2.5mM dNTP mix

25

1.5 µl of primer HWBR-F7 or HWBR/2BPI-475F

1.5 µl of primer HWBR-2BPI-506R or HWBR-1910R

0.75 µl of Expand PCR enzyme (1 unit)

33.25 µl water and

1 µl DNA.

30

Reaction conditions: 10 cycles of 92°C-40 sec; 48°C-1 mm; 68°C-3 mm and 30 cycles of 92°C-40 sec; 55°C- 40 sec; 68°C-2 mm.

Cat. no. V855-20). One clone, named pc3.1/HuGL2R/MH6 (pHuMH6), had the 1.7 kb insert in correct orientation as checked by PCR using vector vs. insert primers.

Functional assay:

5 This hybrid clone was compared to rat GLP-2R using the assay described in Example 2. Results showed that the 2 bp "GA" replacement into the putative deletion site yielded a clone encoding a functional GLP-2R protein, as shown by the CAMP response to GLP-2 treatment.

---

## 10 **EXAMPLE 6**

### **Isolation of the Full-Length Human GLP-2 Receptor cDNA**

Twenty thousand clones from  $\pi$ gt10 cDNA Library from Human Stomach (Clontech; Cat. HL3O17a) were plated on each of 100 agar 150 mm plates. SM buffer (0.1 M NaCl, 10 mM  $Mg_2SO_4$ , 35 mM Tris, pH-7.5, 0.01% gelatin) was added to each plate to  
15 obtain 100 phage lysates each containing 20,000 (20K) pooled clones. The first fifty 20K phage lysates (20K pools) were screened by PCR using two primers designed from HHT13 DNA sequence. The template DNA from each pool was prepared by boiling phage lysate for 10 minutes and centrifuging for 10 minutes.

HWBR-113F [5'-GTGGAGAGGATTTGTGCAAACATTTTC-3'] (SEQ ID NO:24)

20 HWBR-578R [5'-AGAGACATTTCCAGGAGAAGAATGAG-3'] (SEQ ID NO:25)

1  $\mu$ l of each 20K pool DNA was diagnosed by PCR with HWBR-113F and HWBR578R primers using the following conditions:

2  $\mu$ l of 10x Expand™ Buffer 1

2.8  $\mu$ l of 2.5mM dNTP mix

25 0.6  $\mu$ l of primer HWBR-113F

0.6  $\mu$ l of primer HWBR-578R

0.3  $\mu$ l of Expand PCR enzyme (1 unit)

12.7  $\mu$ l water

1  $\mu$ l 20K pool DNA

30 Reaction conditions: 35 cycles of 92°C-40s.; 60°C-40s.; 68°C-1 min.

A 700 bp fragment obtained by Kpn I and Pvu II restriction digest of the amplified DNA from clone UST 38-4-30, and 1.4 kb DNA fragment from Xho I and Pvu II restricted pHHT13-1 DNA were subcloned into Kpn I and Xho I restricted pcDNA3 in a three-way ligation. The new plasmid construct was called pc3/HuGL2R-2. In this manner the full length sequence of the human GLP-2 receptor was obtained.

#### Functional Assay:

The new clone was compared to the rat GLP-2R clone 587 as described previously above. Results showed that the clone encoded a functional human GLP-2R protein, which led to cAMP production in COS cells in response to GLP-2 treatment (Figure 8).

### **EXAMPLE 8 Antibodies directed to the GLP-2 receptor**

#### **1. Antipeptide antibodies**

Antipeptide antibodies were raised in rabbits against an N-terminal peptide (QTRENTTDIWQDESE) (SEQ ID NO:26), a C-terminal peptide (SEG DGSETLQKLR) (SEQ ID NO:27) and extracellular loop 1 (SHNSUSKRPDDESG) (SEQ ID NO:28) of the rat GLP-2 receptor.

Immunocytochemical analysis of serum produced as above confirmed that serum contained antibodies directed to the GLP-2 receptor which do not cross react with the rat GLP-1 receptor.

#### **2. Antibodies to a GLP-2 receptor raised against a fusion protein**

Polynucleotide encoding the C-terminal region of rat GLP-2 receptor (amino acids 444-550) was spliced to the C-terminus of glutathione S-transferase (GST) in pGEX-2T and expressed in E.coli strain SURE1. Protein was purified using affinity chromatography using the above GLP-2 C-terminal fragment fused to the C-terminus of maltose binding protein. Protease degradation was minimized by using a cocktail of protease inhibitors (Boeringer Mannheim).

Antibodies were raised generally according to the method disclosed in Antibodies: A laboratory manual, Harlow and Lane, Cold Spring Harbor Laboratory, 1988. Briefly, the

What is claimed is:

1. An isolated polynucleotide encoding a GLP-2 receptor.
- 5 2. The isolated polynucleotide according to claim 1, which codes for the human GLP-2 receptor having the amino acid sequence of amino acids 67-553 of SEQ ID NO:12
3. The isolated polynucleotide according to claim 2, comprising nucleotides 318-1780 of SEQ ID NO:11
- 10 4. The isolated polynucleotide according to claim 1, which codes for the rat GLP-2 receptor having the amino acid sequence of amino acid 67-550 of SEQ ID NO:2
- 15 5. The isolated polynucleotide according to claim 4, having nucleotides 335-1789 of SEQ ID NO: 1.
6. An isolated polynucleotide which hybridizes under conditions of high stringency with a polynucleotide defined in claim 3 or claim 5.
- 20 7. An oligonucleotide comprising at least 15 nucleotides and having a polynucleotide sequence that hybridizes selectively to a polynucleotide according to any of claims 1-6.
8. An oligonucleotide according to claim 7, wherein the nucleotide sequence of the oligonucleotide corresponds identically with a region of the polynucleotide defined in claim 3 or claim 5.
- 25 9. An oligonucleotide according to claim 7, wherein the nucleotide sequence of the oligonucleotide corresponds identically with a region common to the polynucleotides defined in claim 3 and claim 5.

30

10. In labelled form, a polynucleotide selected from a polynucleotide as defined in any of claims 1-6, and an oligonucleotide as defined in any of claims 7-9.

5 11. A recombinant polynucleotide comprising a GLP-2 receptor-encoding polynucleotide as defined in any of claims 1- 6, and expression controlling elements linked operably therewith to drive expression thereof.

12. A cell that has been genetically engineered by the incorporation expressibly therein of a polynucleotide according to any of claims 1-6.

10 13. The cell according to claim 11, which is a mammalian cell.

14. An antibody which selectively binds to a GLP-2 receptor.

15 15. A recombinant GLP-2 receptor.

16. The GLP-2 receptor of claim 15, comprising amino acids 67-553 of SEQ ID NO:2.

17. The GLP-2 receptor of claim 15, comprising amino acids 67-550 of SEQ ID NO: 12

20 18. A method for identifying GLP-2 receptor ligands, which comprises the steps of:

25 (1) incubating a candidate ligand with a cell as defined in claim 12 or with a membrane preparation obtained therefrom; and then

(2) determining whether binding between the GLP 2 receptor and the candidate ligand has occurred.

30 19. A GLP-2 receptor ligand identified by the method of claim 18.

1/13

1 AAGCTTCGCG GCCTCTGCAG AKGAGTTGTG CAAACACTTC CTCTCTGGAC  
51 AAGGAGGAAT GCAGGAGGCC ACCGCCTGCA GTACATCTTG GAGTGTTGGA  
101 GGGATGTGCC TGCATTGTG AACGGGCGCC AGGAGAATGA GGCCCAACC  
151 AAGCCCGGCA GTGCCAGTA GATGCAGAGA GGCACCCGTG CCCCAGTGTA  
201 GGGCACAGCC AGTGGGCATC CCTGAGGCC AGGGGCCCGT TCCTCTCCAC  
251 TCCCAACAGA TCGTCTGCT GTGGGGCCCT GGGAGGCCCT TCCTCGCCCT  
301 GCTTCTGCTG GTTTCATCA AGCAAGTTAC AGGATCGCTC CTCAAGGAGA  
351 CAACTCAGAA GTGGGCTAAT TATAAGGAGA AGTGTCTGGA AGACTTGCAC  
401 AATAGACTTT CTGGCATATT TTGTAATGGG ACATTTGATC GGTATGTGTG  
451 CTGGCCTCAT TCTTATCCTG GAAATGTCTC TGTTCCCTGT CCTTCATACT  
501 TACCTTGGTG GAATGCAGAG AGCCCAGGAA GGGCCTACAG ACACTGCTTG  
551 GCTCAGGGGA CTTGGCAGAC GCGAGAGAAC ACCACAGATA TTTGGCAGGA  
601 TGAATCAGAA TGCTCAGAGA ACCACAGCTT CAGACAAAAC GTGGATCACT  
651 ACGCCTTGCT ATACACCTTG CAGCTGATGT ACACTGTGGG CTACTCCGTG  
701 TCTCTCATCT CCCTCTTCTT GGCTCTTACA CTCTTCTTGT TCCTTCGAAA  
751 ACTGCATTGC ACACGCAATT ACATCCACAT GAACCTGTTT GCTTCGTTCA  
801 TCCTGAAAGT TCTGGCTGTC CTGGTGAAGG ACATGGTCTC CCACAACTCT  
851 TACTCCAAGA GGCCCGATGA TGAGAGTGGA TGGATGTCAT ATCTGTCAGA  
901 GACATCCGTC TCCTGTGCT CCGTCCAGGT CCTCCTGCAC TACTTTGTGG  
951 GCACCAATCA CTTGTGGCTG CTGGTTGAAG GACTTTACCT CCACACTCTG  
1001 CTGGAGCCCA CAGTGTTTCC TGAAAGGCGG CTGTGGCCCA AGTACCTGGT  
1051 GGTGGGTTGG GCCTTCCCA TGCTGTTTGT TATTCCTGG GGTTTTGCCC  
1101 GTGCACACCT GGAGAACACA CGGTGCTGGG CCACAAATGG GAACCTGAAA  
1151 ATCTGGTGGA TCATCAGAGG ACCCATGCTG CTTTGTGTAA CAGTTAATTT  
1201 CTTTCATCTT CTCAAGATTC TCAAGCTTCT CATTTCTAAG CTCAAAGCTC  
1251 ATCAGATGTG CTTTCAGAGAC TACAAATACA GATTGGCGAA ATCAACGTTG  
1301 CTCCTCATTC CTTTGTGGG GTTTCATGAG GTCCTCTTCA CTTTCTTCCC  
1351 CGACGACCAA GTTCAAGGAT TTTCAAAACG TATTCGACTC TTCATCCAGC  
1401 TGACACTGAG CTCTGTCCAC GGATTTCTGG TGGCCTTGCA GTATGGCTTT  
1451 GCCAATGGAG AGGTGAAGGC AGAGCTGCGA AAGTCATGGG GCCGCTTCTT  
1501 ATTAGCCCGC CACTGGGGCT GCAGAACCTG TGTCTGGGG AAGAATTTCC  
1551 GGTTCTTGGG GAAGTGTTCC AAGAAGCTGT CGGAGGGAGA TGGCTCTGAG  
1601 ACACTCCAGA AGCTGCGGTT CTCCACATGC AGCTCACACC TGGCCTCTGA  
1651 GACCCTGGGA GACGTTGGGG TACAGCCTCA CAGGGGCCGT GGAGCTTGGC  
1701 CCCGGGGAAG CAGCCTGTCT GAGAGCAGTG AGGGAGACTT CACCCTGGCC  
1751 AATACGATGG AGGAGATTCT GGAAGAGAGT GAGATCTAAG GCAGGTCCA

FIG.1A

SUBSTITUTE SHEET

2/13

1801	TCACCGCAGC	TTGGCCACTG	ARGAmCCAAC	CcTARGAAGG	ATKTTGCCGA
1851	RCCCARGGTC	CTCCTCTTCC	TATGTWCtAT	MCCCATTTTG	ATGTGAAGTC
1901	TCTCCTGGGT	GAMCAASCTC	TGTACCAACS	ARTCTCAgTC	CCTCTTGCCC
1951	TTGTCAcCCT	ACTACCCCTC	CCCCATCαCα	CATgTTTTCC	AGAATKTCCG
2001	TTGGTTTGGG	GGGGGGGGTC	TTGCCCTAAA	TTCAAGTsGA	GTGGARCCCA
2051	CCATGAAGAA	AARTCATTTA	TTAAATAGAR	TCCGGTTAGG	ATCTCCTTCC
2101	CGTTCATGGT	GCATGGCCTC	CTTCCAAGGG	ATGGGAGTCG	GSTGCACTGG
2151	AACCCACAG	GAAAcTTTGA	AGTATCCAGT	TCTAGGGAAT	TATAGCCAAT
2201	ATTCTGAGAG	AGCAAGTCTG	AGATGAGAKC	CGAGAATWGC	AAGTGTWGGA
2251	CAWGCATTCA	AGGAAACTCC	TCACCTTTgG	GCGAAACCTA	tGGCAGGATc
2301	GGCAtGGAGC	AGCTATTMTG	CAAYGGCCGC	TCACCTGGGA	CATACCACTC
2351	TCCTTGGGCA	GGATGTGACC	CCATGTKGTC	CCCCAGACTC	CTCTCCTCCT
2401	TGCTTSTSTT	CYTTTCCYGT	CAAGTCTCAC	CTCCCTTTCT	ACATCTCAGT
2451	TCWGTTTGGT	GTYGACAGAA	GYTGAATGT	CACAATACTG	CATGTGTTAG
2501	TTTCTGTCGT	CATTGCTGTG	TCCAAATACC	TGACCAGGAC	CAATTTAAGC
2551	GAGGAACTGC	TACATGGGCG	GCCGC		

FIG.1B

SUBSTITUTE SHEET

20 APRIL 1998 (20.04.98)

3/13

1 MRPQPSPAVP SRCREAPVPR VRAQPVGIPE AQGPVPLHSQ QMRLLWGPGR  
51 PFLALLLLVS IKQVTGSLK ETTQKWANYK EKCLEDLHNR LSGIFCNGTF  
101 DRYVCWPHSY PGNVSVPCPS YLPWWNAESP GRAYRHCLAQ GTWQTRENTT  
151 DIWQDESECS ENHSFRQNV D HYALLYTLQL MYTVGYSVSL ISLFLALTLF  
201 LFLRKLHCTR NYIHMNLFAS FILKVLAVLV KDMVSHNSYS KRPDDESGWM  

---

251 SYLSETSVSC RSVQVLLHYF VGTNHLWLLV EGLYLHTLLE PTVPERRRLW  
301 PKYLVVGWAF PMLFVIPWGF ARAHLENTRC WATNGNLKIW WIIRGPMLLC  
351 VTVNFFIFLK ILKLLISKLK AHQMCFRDYK YRLAKSTLLL IPLLGVHEVL  
401 FTFFPDDQVQ GFSKRIRLFI QLTLSVHGF LVALQYGFAN GEVKAELRKS  
451 WGRFLLARHW GCRTCVLGKN FRFLGKCSKK LSEGDCSETL QKLRFCSTCSS  
501 HLASETLGDV GVQPHRGRGA WPRGSSLSSES SEGDFTLANT MEEILEESEI

FIG.2

SUBSTITUTE SHEET



4/13




FIG.3

SUBSTITUTE SHEET

20 APRIL 1998 (20.04.98)

5/13

1 TCCTTCTCTC TTATCTCCCT CTTCTGGCT CTGACCCTCC TCTTGTTTCT  
51 TCGAAAACCTC CACTGCACGC GCAACTACAT CCACATGAAC TTGTTTGCTT  
101 CTTTCATCCT GAGAACCCTG GCTGTACTGG TGAAGGACGT CGTCTTCTAC  
151 AACTCTTACT CCAAGAGGCC TGACAATGAG AATGGGTGGA TGTCTTACCT  
201 GTCAGAGATG TCCACCTCCT GCCGCTCAGT CCAGGTTCTC TTGCATTACT  
251 TTGTGGGTGC CAATTACTTA TGGCTGCTGG TTGAAGGCCT CTACCTCCAC  
301 ACGCTGCTGG AGCCACAGT GCTTCCTGAG AGGCGGCTGT GGCCCARATA  
351 CCTGCTGTTG GGTGGGCCT TCCCTGTGCT ATTTGTTGTA CCCTGGGGTT  
401 TCGCCCGTGC ACACCTGGAR AACACAGGGT GCTGGACAAC AAATGGGAAT  
451 AAGAAAATCT GGTGGATCAT CCGAGGACCC ATGATGCTCT GTGTAACAGT  
501 CAATTTCTTC ATCTTCCTGA AAATTCTCAA GCTTGTCATT TCTAAGCTCA  
551 AAGCTCATCA AATGTGCTTC AGAGATTATA AATACAGATT GGCAAAATCA  
601 AACTGGTCC TCATTCCTTT ATTGGGCGTT CATGAGATCC TCTTCTCTTT  
651 CATCACTGAT GATCAAG

FIG.4

SUBSTITUTE SHEET

1	S F S L I S L F L A L T L L L F L R K L TCCTTCTCTCTTATCTCCCTCTTCCTGGCTCTCACCCCTCTTGTTCCTTCGAAAACCTC	80
61	H C T R N Y I K M N L F A S F I L R T L CACTGCACGCGCAACTACATCCACATGAACCTGTTTGCTTCTTTTCATCCTGAGAACCCTG	120
121	A V L V K D V V F Y N S Y S K R P D N E GCTGTACTGGTGAAGGACGTCGTCTTCTACAACTCTTACTCCAAGAGGCCTGACAATGAG	180
181	N G W M S Y L S E M S T S C A S V Q V L AATGGGTGGATGTCTACCTGTCAGAGATGTCCACCTCCTGCCGCTGAGTCCAGGTTGTG	240
211	L H Y F V G A N Y L W L L V E G L Y L H TTGCATTACTTTGTGGGTGCCAATTACTTATGGCTGCTGGTTGAAGGCCTCTACCTCCAC	300
301	T L L E P T V L P E R R L W P K Y L L L ACGCTGCTGGAGCCACAGTGCTTCCTGAGAGGCGGCTGTGGCCCA?ATACCTGCTGTTG	360
361	G W A F F V L F V V P W G F A R A H L E GGTTGGGCCTTCCCTGTGCTATTTGTTGTACCCTGGGGTTTTCGCCCGTGACACCTGGAR	420
421	N T G C W T F N G N K K I W W I I R G P AACACAGGGTGCTGGACAACAAATGGGAATAAGAAAATCTGGTGGATCATCCGAGGACCC	480
481	M M L C V T V N F F I F L K I L K L L I ATGATGCTCTGTGTAACAGTCAATTTCTTCATCTTCTGAAAATTCTCAAGCTTCTCATT	540
541	S K L K A H Q M C F R D Y K Y R L A K S TCTAAGCTCAAAGCTCATCAATGTGCTTCAGAGATTATAAATACAGATTGGCAAAATCA	500
501	T L V L I P L L G V H E I L F S F I F D ACACTGGTCCTCATTCCTTTATTGGGCGTTCATGAGATCCTCTTCTCTTTCATCACTGAT	560
661	D Q GATCAAG	667

7/13

1	TGGAGAGGATTTGTGCAACATTTCTTCTGTGGACCAAGAGGAATGCAAGAGGAGGCTGC	60
61	CTGCGGTGCATCTTGGACGGCTAGAGAGATGTACCCCTACTTGTGAAGGTGCACGAGGAA	120
121	M K L G S S R A G P G R G S A G L L P G GATGAAGCTGGGATCGAGCAGGGCAGGGCCTGGGAGAGGAAGCGCGGGACTCCTGCCTGG	180
181	V H E L P M G I P A P W G T S P L S F H CGTCCACGAGCTGCCCATGGGCATCCCTGCCCCCTGGGGGACCAGTCCTCTCTCTTCCA	240
241	R K C S L W A P G R P F L T L V L L V S CAGGAAGTGCTCTCTCTGGGCCCTGGGAGGCCCTTCTCACTCTGGTCCTGCTGGTTTC	300
301	I K Q V T G S L L E E T T R K W A Q Y K CATCAAGCAAGTTACAGGATCCCTCGTTGAGGAAACGACTCGGAAGTGGGCTCAGTACAA	360
361	Q A C L R D L L K E P S G I F C N G T F ACAGGCATGTCTGAGAGACTTACTCAAGGAACCTTCTGGCATATTTGTAACGGGACATT	420
421	D Q Y V C W P H S S P G N V S V P C P S TGATCAGTACGTGTGTGGCCTCATTCTTCTCTGGAAATGTCTCTGTACCCTGCCCTTC	480
480	Y L P W W S E E S S G R A Y R H C L A Q ATACTTACCTTGGTGGAGTGAAGAGAGCTCAGGAAGGGCTACAGACACTGCTTGGCTCA	540
541	G T W Q T I E N A T D I W Q D D S E C S GGGGACTTGGCAGACGATAGAGAACGCCACGGATATTTGGCAGGATGACTCCGAATGCTC	600
601	E N H S F K Q N V D R Y A L L S T L Q L CGAGAACCACAGCTTCAAGCAAAACGTGGACCGTTATGCCTTGCTGTCAACCTTGCAGCT	660

FIG.6A

SUBSTITUTE SHEET

8/13

M Y T V G Y S F S L I S L F L A L T L L  
661 GATGTACACCGTGGGATACTCCTTCTCTCTTATCTCCCTCTTCTGGCTCTCACCTCCT  
-----+-----+-----+-----+-----+-----+-----+ 720

L F L R K L H C T R N Y I H M N L F A S  
721 CTGTTTCTTCGAAACTCCACTGCACGCGCAACTACATCCACATGAACCTGTTTGCTTC  
-----+-----+-----+-----+-----+-----+-----+ 780

---

F I L R T L A V L V K D V V F Y N S Y S  
781 TTTCATCCTGAGAACCCTGGCTGTACTGGTGAAGGACGTCGTCTTCTACAACTCTTACTC  
-----+-----+-----+-----+-----+-----+-----+ 840

K R P D N E N G W M S Y L S E M S T S C  
841 CAAGAGGCCTGACAATGAGAATGGGTGGATGTCTACCTGTCAGAGATGTCCACCTCCTG  
-----+-----+-----+-----+-----+-----+-----+ 900

R S V Q V L L H Y ? V G A N Y L W L L V  
901 CCGCTCAGTCCAGGTTCTCTTGCACTTTGTGGGTGCCAATTACTTATGGCTGCTGGT  
-----+-----+-----+-----+-----+-----+-----+ 960

TGAAGGCCTCTACCTCCACACGCTGCTGGAGCCCACAGTGCTTCCTGAGAGGCGGCTGTG  
961 -----+-----+-----+-----+-----+-----+-----+ 1020

P R Y L L L G W A F P V L F V V P W G F  
1021 GCCCAGATACCTGCTGTTGGGTTGGGCCTTCCCTGTGCTATTTGTTGTACCCTGGGGTTT  
-----+-----+-----+-----+-----+-----+-----+ 1080

A R A H L E N T G C W T T N G N K K I W  
1080 CGCCCGTGCACACCTGGAGAACACAGGGTGCTGGACAACAAATGGGAATAAGAAAATCTG  
-----+-----+-----+-----+-----+-----+-----+ 1140

W I I R G P M M L C V T V N F F I F L K  
1141 GTGGATCATCCGAGGACCCATGATGCTCTGTGTAACAGTCAATTTCTTCATCTTCCTGAA  
-----+-----+-----+-----+-----+-----+-----+ 1200

I L K L L I S K L K A H Q M C F R D Y K  
1201 AATTCTAAGCTTCTCATTCTAAGCTCAAAGCTCATCAATGTGCTTCAGAGATTATAA  
-----+-----+-----+-----+-----+-----+-----+ 1250

Y R L A K S T L V L I P L L G V H E I L  
1261 ATACAGATTGGCAAAATCAACACTGGTCCTCATTCCTTTATTGGGCGTTCATGAGATCCT  
-----+-----+-----+-----+-----+-----+-----+ 1320

F S F I T D D Q V E G F A K L I R L F I  
1321 CTTCTCTTTCATCACTGATGATCAAGTTGAAGGATTGCAAACTTATACGACTTTTCAT  
-----+-----+-----+-----+-----+-----+-----+ 1380

FIG. 6B  
SUBSTITUTE SHEET

20 APRIL 1998 (20.04.98)

9/13

Q L T L S S F H G F L V A L Q Y G F A N  
TCAGTTGACACTGAGCTCCTTTTCATGGGTTCTGGTGGCCTTGACAGTATGGTTTTGCCAA  
1381 -----+-----+-----+-----+-----+-----+-----+ 1440

G E V K A E L R K Y W V R F L L A R H S  
TGGAGAAGTGAAGGCTGAGCTGCCGAAATACTGGGTCCGCTTCTTGCTAGCCCGCCACTC  
1441 -----+-----+-----+-----+-----+-----+-----+ 1500

G C R A C V L G K D F R F L G K C P K K  
AGGCTGCAGAGCCTGTGTCCTGGGGAAGGACTTCCGGTTCCTAGGAAAATGTCCCAAGAA  
1501 -----+-----+-----+-----+-----+-----+-----+ 1560

L S E G D G A E K L R K L Q P S L N S G  
GCTCTCGGAAGGAGATGGCGCTGAGAAGCTTCGGAAGCTGCAGCCCTCACTTAACAGTGG  
1561 -----+-----+-----+-----+-----+-----+-----+ 1620

R L L H L A M R G L G E L G A Q P Q Q D  
GCGGCTCCTACATCTAGCCATGCGAGGTCTTGGGGAGCTGGGCGCCAGCCCCAACAGGA  
1621 -----+-----+-----+-----+-----+-----+-----+ 1680

H A R W P R G S S L S E C S E G D V T M  
CCATGCACGCTGGCCCCGGGGCAGCAGCCTGTCCGAGTGCAGTGAGGGGGATGTCACCAT  
1681 -----+-----+-----+-----+-----+-----+-----+ 1740

A N T M E E I L E E S E I  
GGCCAACACCATGGAGGAGATTCTGGAAGAGAGTGAGATCTAGGGTGGAGTTCCACCACC  
1741 -----+-----+-----+-----+-----+-----+-----+ 1800

CTGGCTCTGCTCCCAGGGACTCTTGAGGGGGGCCAGGAAGAGGAAGCAAAGCAGGACACA  
1801 -----+-----+-----+-----+-----+-----+-----+ 1860

CGTTGCTGGGCACGGAATCATTCTCGTTCCATTACCATGCCACTTTGATATGAAAGCTA  
1861 -----+-----+-----+-----+-----+-----+-----+ 1920

TCACAAGGTTCTTCAAGCTCTGTATGAAAGAGGCTGTGTGTCATGCTCACAGCCTCTGCC  
1921 -----+-----+-----+-----+-----+-----+-----+ 1980

TGCTCTTCTCATCCTAATAACCCCCACCAGTGTGTTTTCCACAATGCCACCAGACCCTA  
1981 -----+-----+-----+-----+-----+-----+-----+ 2040

GGGCCTGGCTCTAAATTCAAGCCAATGAAGTCCCACCCGGAATTCTTTTGCTTTTACCC  
2041 -----+-----+-----+-----+-----+-----+-----+ 2100

CTGGAAGAAATA  
2101 -----+----- 2112

FIG.6C  
SUBSTITUTE SHEET

20 APRIL 1998 (20.04.98)

10/13

1 MKLGSSRAGP GRGSAGLLPG VHELPMGIPA PWGTSPLSFH RKCSLWAPGR  
51 PFLTLLVLLVS IKQVTGSLLE ETTRKWAQYK QACLRDLLE PSGIFCNGTF  
101 DQYVCWPHSS PGNVSVPCPS YLPWWSEESS GRAYRHCLAQ GTWQTIENTAT  
151 DIWQDDSECS ENHSFKQNV DRYALLSTLQL MYTVGYFSFSL ISLFLALILL  
201 LFLRKLHCTR NYIHMNLFAS FILRTLAVLV KDVFYNSYS KRPDNENGWM  
251 SYLSEMSTSC RSVQVLLHYF VGANYLWLLV EGLYLHTLLE PTVLPERRLW  
301 ~~PRYLLLGWAF PVLFVVPWGF ARAHLENTGC WTTNGNKKIW WIIRGPMMLC~~  
351 VTVNFFIFLK ILKLLISKLK AHQMCFRDYK YRLAKSTLVL IPLLGVHEIL  
401 FSFITDDQVE GFAKLIRLFI QLTLSFHFGL LVALQYGFAN GEVKAELRKY  
451 WVRFLARHS GCRACVLGKD FRFLGKCPKK LSEGDAEKL RKLQPSLNSG  
501 RLLHLAMRGL GELGAQPQQD HARWPRGSSL SECSEGDVTM ANTMEEILEE  
551 SEI

FIG.7

SUBSTITUTE SHEET

11/13




FIG.8



20 APRIL 1998 (20.04.98)

12/13

1	MRPQSPAVPSRCREAPVPRVRAQPVGIP EAQGPVPLHSQQMRLWGPGR	50
	:       .     .	
1	MKLGS SRAGPGRGSAGLLPGVHELPMGIPAPWGTSPLSFHRKCSLWAPGR	50
51	PFLALLLVSIKQVTGSLLKETTKWANYKEKCLEDLHNRLSGIFCNGTF	100
	.	
51	PFLTLLVSIKQVTGSLL EETTRKWAQYKQACLRDLLKEPSGIFCNGTF	100
101	DRYVCWPHSYPGNVSVPCPSYLPWWNAESPGRAYRHCLAQGTWQTRENTT	150
	.	
101	DQYVCWPHSSPGNVSVPCPSYLPWWSESSGRAYRHCLAQGTWQTIENAT	150
151	DIWQDESECSNHSFRQNV DHYALLYLQMYTVGYSVSLISLFLALTLF	200
	:	
151	DIWQDSECSNHSFKQNVDRYALLSTLQMYTVGYSFSLISLFLALTLL	200
201	LFLRKLHCTRNYIHMNLFASFILKVLAVLVKDMVSHNSYSKRPDDESGWM	250
201	LFLRKLHCTRNYIHMNLFASFILRTLAVLVKDVVFYNSYSKRPDNENGWM	250
251	SYLSETSVSCRSVQVLLHYFVG TNHLWLLVEGLYLHTLLEPTVFPERRLW	300
251	SYLSEMSTSCRSVQVLLHYFVGANYLWLLVEGLYLHTLLEPTVLPERRLW	300
301	PKYL VVGWAF PMLFVIPWGFARAHLENTRCWATNGNLKIWWIIRGPMMLC	350
	:     .       .     :	
301	PRYLLL GWAF PVLFVVPWGFARAHLENTGCWTNGNKKIWWIIRGPMMLC	350
351	VTVNFFIFLKILKLLISKLKAHQMCFRDYKYRLAKSTLLLIPLLGVHEVL	400
351	VTVNFFIFLKILKLLISKLKAHQMCFRDYKYRLAKSTLVLP LLGVHEVL	400
401	F TFFPDDQVQGF SKRIRLF IQLTLSSVHGFLVALQYGFANGEVKAELRKS	450
	.       :    .	
401	F SFI TDDQVEGFAKLIRLF IQLTLSSFHGFLVALQYGFANGEVKAELRKY	450
451	WGRFLLARHWGCRTCVLGKNFRFLGKCSKKLSEGDGSETLQKLRFSTCSS	500
451	WVRFLLARHSGCRACVLGKDFRFLGKCPKKLSEGDGAELRKLQPSLNSG	500
501	...HLASETLGDVGVP HRGRGAWPRGSSLSSESSEGDFTLANTMEEILEE	547
	:	
501	RLHLAMRGLGELGAQPQQDHARWPRGSSLSECSEGDVTMANTMEEILEE	550
548	SEI 550	
551	SEI 553	

FIG.9

SUBSTITUTE SHEET

13/13

GL2R\_RAT\_R  
HWBRPAT\_TR  
GLPR\_HUMAN

MRPQSPAVPSRCREAPVPRVRAQPVG IPEAQQPVPLHSQQMRL LWGPG-RPFLALLLV  
MKLGSSRAGPGRGSAGLLPGVHELPMG IPAPWGTSPLSFHRKCSLWAPG\_RPFLTLVLLV  
-----MAGAPGRLRLALLLGMVGRAGPRP-----  
\* \* \*

GL2R\_RAT\_R  
HWBRPAT\_TR  
GLPR\_HUMAN

SIKQVTGSLLKETQKWANYKEKCLEDLHNRL---SGIFCNGTFDRYVCWPHSYPG-NVS  
SIKQVTGSLLLEETTRKWAQYKQACLRDLLKEP---SGIFCNGTFDRYVCWPHSYPG-NVS  
---QGATVSLWETVQKWREYRQCQRSLTEDPPPATDLFCNRTFDEYACWPDGEPGSFVN  
\* \* \* \* \*

GL2R\_RAT\_R  
HWBRPAT\_TR  
GLPR\_HUMAN

VPCPSYLPWWNAESPGRAYRHCLAQGTWQTRENTTDIWQDESECSENHFRQNVHDYALL  
VPCPSYLPWWSEESSGRAYRHCLAQGTWQTIENATDIWQDDSECSENHFRQNVDRYALL  
VSCPWYLPWASSVPQGHVYRFTAEGWLKQDNSSLPWRDLSECSESKRGERSSPEEQLL  
\* \* \* \* \*

GL2R\_RAT\_R  
HWBRPAT\_TR  
GLPR\_HUMAN

YTLQLMYTVGYSVSLISLFLALTFLFLRKLHCTRNYIHMNLFASFILKVLAVLVKDMVS  
STLQLMYTVGYSFSLISLFLALTLLFLRKLHCTRNYIHMNLFASFILRTLAVLVKDVVF  
F-LYIIYTVGYALSFSALVIASAILLGRHLHCTRNYIHLNLFASFILRALSVFIKDAAL  
\* \* \* \* \*

GL2R\_RAT\_R  
HWBRPAT\_TR  
GLPR\_HUMAN

HNSYSKRPDDESGWMSYLS-ETSVSCRSVQVLLHYFVGINHLLWLLVEGLYLHTLLEPTVF  
YNSYSKRPDNENGWMSYLS-EMSTSCRSVQVLLHYFVGANYLLWLLVEGLYLHTLLEPTVL  
KWMYST-AAQQHQWDGLLSYQDSLSCRLVFLMQYCAANYWLLVEGVLYLTLLAFSVF  
\* \* \* \* \*

GL2R\_RAT\_R  
HWBRPAT\_TR  
GLPR\_HUMAN

PERRLWPKYLVGWAFPMFLVFPWGFARAHLENTRCWATNGNLKIWWIIRGPMMLCVTVN  
PERRLWFRYLLLGWAFPVLFVVPWGFARAHLENTGCWTNGNKKIWWIIRGPMMLCVTVN  
SEQWIFRLYVSIWGVPLLFVVPWGVKYLVEDEGCWTRNSNMNYWLIIRLPILFAIGVN  
\* \* \* \* \*

GL2R\_RAT\_R  
HWBRPAT\_TR  
GLPR\_HUMAN

FFIFLKILKLLISKLAHQMCFRDYKYRLAKSTLLLIPLLGVHEVLFTFFPDDQVQGF SK  
FFIFLKILKLLISKLAHQMCFRDYKYRLAKSTLVLLIPLLGVHEILFSF ITDDQVEGFAK  
FLIFVRVICIVSKLKANLMCKTDIKCRLAKSTLTLIPLLGTHEVIFAFVMDHARGTLR  
\* \* \* \* \*

GL2R\_RAT\_R  
HWBRPAT\_TR  
GLPR\_HUMAN

RIRLF IQLTLSSVHGFLVALQYGFANGEVKAELRKSWGRFLLARHWGCRTCVLGKNFRFL  
LIRLF IQLTLSSFHGFVALQYGFANGEVKABLYKWRFLARHSGCRACVLGRDFRFL  
FIKLFTELSFTSFQGLMVAIFYCFVNNEVQLEFRKSWEWRLEHLHIQRDSSMKP-----  
\* \* \* \* \*

GL2R\_RAT\_R  
HWBRPAT\_TR  
GLPR\_HUMAN

GKCSKKLSEGDGSETLQKLRFSTCSS--HLASETLGDVGVPQPHRGAWPRGSSLSESS  
GKCPKKLSEGDGAELRKQLPSLNSGRLLHLAMRGLGELGAQPQDQDARWPRGSSLSECS  
LKOPTSS-LSGATAGSSMYTATCQASCS-----  
\* \* \*

GL2R\_RAT\_R  
HWBRPAT\_TR  
GLPR\_HUMAN

EGDFTLANTMEEILEESEI  
EGDVTMANTMEEILEESEI  
-----

FIG.10

SUBSTITUTE SHEET



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07K 14/00</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 98/25955</b> <b>(43) International Publication Date:</b> 18 June 1998 (18.06.98)
<b>(21) International Application Number:</b> PCT/CA97/00969 <b>(22) International Filing Date:</b> 15 December 1997 (15.12.97) <b>(30) Priority Data:</b> 08/767,224 13 December 1996 (13.12.96) US 08/787,721 24 January 1997 (24.01.97) US 08/845,546 24 April 1997 (24.04.97) US <b>(71) Applicant (for all designated States except US):</b> ALLELIX BIOPHARMACEUTICALS INC. [CA/CA]; 6850 Goreway Drive, Mississauga, Ontario L4V 1V7 (CA). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> MUNROE, Donald, G. [CA/CA]; 27 Wakefield Lane, Waterdown, Ontario L0R 2H3 (CA). GUPTA, Ashwani, K. [CA/CA]; 7031 Dunrobin Way, Mississauga, Ontario L5N 6Y4 (CA). VYAS, Tejal, B. [CA/CA]; 275 Riel Drive, Mississauga, Ontario L5B 3K1 (CA). McCALLUM, Kirk [CA/CA]; 3918 Pitch Pine Crescent, Mississauga, Ontario L5L 1P8 (CA). FAN, Ermi [CA/CA]; Apartment #418, 20 Southport Street, Toronto, Ontario M6S 4Y8 (CA). <b>(74) Agent:</b> AITKEN, David, W.; Osler, Hoskin & Harcourt, Suite 1500, 50 O'Connor Street, Ottawa, Ontario K1P 6L2 (CA).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> CLONED GLUCAGON-LIKE PEPTIDE-2 RECEPTORS		
<b>(57) Abstract</b>  The invention relates to nucleotides and amino acid sequences encoding glucagon-like peptide 2 receptors, recombinant host cells transformed with such nucleotides, and methods of using the same in drug screening and related applications.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## CLONED GLUCAGON-LIKE PEPTIDE-2 RECEPTORS

### FIELD OF THE INVENTION

The invention is in the field of molecular biology. It relates, more particularly, to cloned glucagon-like peptide 2 receptors and their use in drug screening and related applications.

### BACKGROUND TO THE INVENTION

Glucagon-like peptide-2 (GLP-2) is a 33 amino acid peptide, which is expressed in a tissue determined fashion from the pleiotrophic glucagon gene and is highly related in terms of amino acid sequence to glucagon and Glucagon-like peptide-1 (GLP-1). Mammalian forms of GLP-2 are highly conserved: for example, the human and degu (a south American rodent) forms differ by one and three amino acids respectively from rat GLP-2. Recently it was demonstrated that GLP-2 is an intestinotrophic peptide hormone; when given exogenously, GLP-2 can produce a marked increase in the proliferation of small intestinal epithelium of the test mice (Drucker et al, (1996) PNAS, 93:7911-7961). More recently, GLP-2 has been shown to increase D-Glucose maximal transport rate across the intestinal basolateral membrane (Cheeseman and Tseng: American Journal of Physiology (1996) 271:G477-G482).

To accelerate research into gastrointestinal biology and development of drugs useful in the treatment of various medical conditions including gastrointestinal disorders, it would be useful to provide the receptor through which the effects of GLP-2 are mediated.

### SUMMARY OF THE INVENTION

The GLP-2 receptor has now been cloned and characterized. Accordingly, the present invention provides an isolated polynucleotide encoding a GLP-2 receptor, particularly including mammalian forms and homologs thereof such as, in specific embodiments, the rat and human forms. In aspects of the invention, polynucleotide coding for a GLP-2 receptor is utilized for expression to obtain functional receptor protein and, in optionally labelled form, for further gene cloning to identify structurally related receptor proteins. In related aspects of the invention, anti-sense versions of GLP-2 receptor-encoding polynucleotides and fragments thereof are obtained and utilized to regulate GLP-2 receptor expression.

In another of its aspects, the invention provides GLP-2 receptor as a product of recombinant production in a cellular host. In related aspects, there are provided recombinant host cells that express GLP-2 receptor, as well as receptor-bearing membranes derived from such cells, and expression constructs in which polynucleotide coding for the GLP-2 receptor is linked to expression controls functional in the selected host cell.

In another of its aspects, the GLP-2 receptor is utilized in a chemicals screening program to identify GLP-2 receptor ligands. This method comprises the steps of incubating the candidate ligand with a GLP-2 receptor-producing cell of the present invention, or with a membrane preparation derived therefrom, and then measuring whether, or the extent to which, binding has occurred. Using cells that express a GLP-2 receptor coupled functionally to a second messenger system, such binding can be determined indirectly, to reveal ligand against activity, by detecting an appropriate reporter.

In another of its aspects, the invention provides antibodies directed to the GLP-2 receptor, for use for example in diagnostic procedures.

The invention is further described with reference to the following drawings in which:

#### BRIEF REFERENCE TO THE DRAWINGS

Figure 1 discloses a cDNA sequence (SEQ ID NO: 1), nucleotides 137-1789 of which encode the rat GLP-2 receptor wherein ambiguous base pairs are indicated using the standard IUB nomenclature (R: A or G, Y: C or T, M: A or C, K: G or T, S: G or C, W: A or T).

Figure 2 discloses the amino acid sequence of the expression product (SEQ ID NO: 2) from the cDNA of Figure 1.

Figure 3 illustrates the relative potencies of GLP-2 peptide and GLP-1 peptides for the receptor encoded by SEQ ID NO: 1.

Figure 4 discloses a cDNA sequence of 667 nucleotides (SEQ ID NO: 9) which encodes a 222 amino acid fragment (SEQ ID NO: 10) of a human GLP-2 receptor.

Figure 5 discloses the amino acid sequence (SEQ ID NO: 10) expressed from the cDNA of Figure 4.

Figure 6 discloses a cDNA sequence (SEQ ID NO: 11), nucleotides 121-1779 of which encode a human GLP-2 receptor.

Figure 7 discloses the amino acid sequence of the expression product (SEQ ID NO: 12) from the cDNA of Figure 6.

Figure 8 illustrates the functional activation by GLP-2 peptide of the human receptor encoded by SEQ ID NO: 11 (Figure 8).

Figure 9 compares the amino acid sequences of the rat GLP-2 receptor and the human GLP-2 receptor.

Figure 10 compares the amino acid sequences of the rat GLP-2 receptor and the human GLP-2 receptor against rat GLP-1 receptor.

#### DETAILED DESCRIPTION AND PREFERRED EMBODIMENTS

The invention relates in one respect to polynucleotides, in their isolated form, that code for GLP-2 receptors. As used herein "isolated" means separated from polynucleotides that encode other proteins. In the context of polynucleotide libraries, for instance, GLP-2 receptor-encoding polynucleotide is considered "isolated" when it has been selected, and hence removed from association with other polynucleotides within the library. Such polynucleotides may be in the form of RNA, or in the form of DNA including cDNA, genomic DNA and synthetic DNA. The GLP-2 receptors are characterized by structural features common to the G-protein coupled receptor class, including seven transmembrane regions, and by the functional properties of binding GLP-2 peptide selectively, i.e.,

preferentially, to GLP-1 peptide. Such selective binding is revealed as a statistically significant greater binding affinity of GLP-2 than of GLP-1, in the context of the assay chosen to measure such affinity. When expressed functionally in a host cell, i.e., in operable linkage with a responsive second messenger system the GLP-2 receptors are capable further of responding to GLP-2 binding by signal transduction. In this regard, the activity of a G-protein coupled receptor such as a GLP-2 receptor can be measured using any of a variety of appropriate functional assays in which activation of the receptor results in detectable change in the level of some second messenger system, such as adenylyl cyclase, calcium mobilization, inositol phospholipid hydrolysis products or guanylyl cyclase.

With reference to Figure 9 and Figure 10, which reveals homologies across amino acid sequences representing human and rat GLP-2 receptors, regions of 100% identity are indicated by solid vertical bars. In embodiments of the invention, the GLP-2 receptors are defined structurally as receptors that incorporate these regions of amino acid sequence and that also exhibit the functional characteristic of binding GLP-2 peptide selectively, relative to GLP-1 peptide. In more specific embodiments, the GLP-2 receptor structure further incorporates those amino acids which, across the human and rat receptor species, are highly conserved (indicated by ':'). At these sites, it will be appreciated that the sequence can contain any amino acid within the highly conserved amino acid family to which the identified amino acid belongs. In still more specific embodiments, the GLP-2 receptors have a structure which still further incorporates to the moderately conserved amino acids (indicated by '.') meaning, at these sites, that the amino acids within the moderately conserved family to which they belong. Beyond these sequences, the GLP-2 receptor structure can vary widely in embodiments of the invention, in allowing for non-conservative amino acid substitutions.

In one embodiment of the invention, the GLP-2 receptor is a rat GLP-2 receptor having the amino acid sequence of SEQ ID NO:2. In a specific embodiment of the invention, this rat GLP-2 receptor is encoded by the polynucleotide sequence of SEQ ID NO: 1. This particular GLP-2 receptor-encoding polynucleotide, also referred to as the WBR



gene, is a cDNA of rat origin. The expression product of this polynucleotide incorporates the mature form of the GLP-2 receptor, and additionally incorporates a secretion signal that is removed before membrane integration of the mature GLP-2 receptor product. Such a signal sequence may be naturally present on the polypeptides or replaced with a functionally equivalent secretion signal which is heterologous to the GLP-2 receptor. The replacement secretion signal chosen will depend on the expression system being used and will typically, but not essentially be endogenous to the chosen host and, also typically but not essentially, be homologous to the chosen expression controlling sequences.

The expressed rat GLP-2 receptor product (Figure 2, SEQ ID NO: 2) is characterized structurally as a single 550 amino acid polypeptide chain having a predicted molecular weight of 72kDa. Two functional translation start sites of the rat GLP-2 receptor have been identified, these are the codons encoding methionine 1 and methionine 42 of SEQ ID NO:2. Without wishing to be limited, by analogy with the GLP-1 receptor, it is believed that residues 1-66 of SEQ ID NO:2 are cleaved to provide a mature protein (i.e., the amino acid sequence of the receptor as it appears in the cell membrane) of 484 amino acids. With respect to structural domains of this GLP-2 receptor, hydropathy analysis and sequence alignment with related members of this sub-family of G protein coupled receptors indicates seven putative transmembrane domains, one spanning residues 181-203 inclusive (TM I), another spanning residues 211-230 (TM II), a third spanning residues 262-285 (TM III), a fourth spanning residues 300-321 (TM VI), a fifth spanning 339-362 (TM V), a sixth spanning 386-405 (TM VI) and a seventh spanning 422-441 (TM VII). Based on this assignment, it is likely that this GLP-2 receptor, in its natural membrane-bound form, consists of a an N-terminal extracellular domain, followed by a hydrophobic region containing seven transmembrane domains and an intracellular 442-550 amino acid C-terminal domain. The protein exhibits the highest degree of homology to the rat GLP-1 receptor with 49% identity at the amino acid level.

In a related embodiment, the GLP-2 receptor is of human origin (SEQ ID NO: 9) and incorporates the human GLP-2 receptor fragment having the amino acids of SEQ ID NO: 10.

This polynucleotide was isolated using the rat cDNA sequence as generally described below and as detailed in Example 3. In a further related embodiment of the invention, the cDNA is of human origin (SEQ ID NO: 11) and encodes the full length human GLP-2 receptor having residues 67-533 of the amino acid sequence of SEQ ID NO: 12. The human GLP-2 receptor precursor product (Figure 7, SEQ ID NO: 12) is characterized structurally as a single 553 amino acid polypeptide chain having a predicted molecular weight of 72 kDa. It is believed that, as for the rat GLP-2 receptor, this sequence, this precursor form of the human GLP-2 receptor incorporates an N-terminal signal sequence, which can be replaced by a functionally equivalent heterologous signal sequence. Without wishing to be limited it is believed that the mature form of the human GLP-2 receptor results after cleavage of residues 1-66 of SEQ ID NO:12 (Figure 7). With respect to structural domains of this GLP-2 receptor, hydropathy analysis and sequence alignment with related members of this sub-family of G protein coupled receptors indicates seven putative transmembrane domains, one spanning residues 181-203 inclusive (TM I), another spanning residues 211-230 (TM II), a third spanning residues 262-285 (TM III), a fourth spanning residues 300-321 (TM IV), a fifth spanning residues 339-362 (TM V), a sixth spanning residues 386-405 (TM VI), and a seventh spanning residues 422-441 (TM VII). Based on this assignment, it is likely that this GLP-2 receptor, in its natural membrane-bound form, consists of an N-terminal extracellular domain, followed by a hydrophobic region containing seven transmembrane domains interspersed with six short hydrophilic domains, and an intracellular domain, which is predicted to span residues 442-553. A second form of this GLP-2 receptor encompassed by the invention has a translational start site at the methionine codon at position 26 of the amino acid sequence presented in Figure 7 SEQ ID No. 12. The resulting 528 amino acid polypeptide chain also consists of an extracellular domain, seven transmembrane domains, and a C-terminal intracellular domain, and is at least 95% identical in sequence to residues 26-553 of the sequence presented in Figure 7, SEQ ID NO: 12.

In another embodiment, the invention provides GLP-2 receptor polynucleotide sequences and their unique sequence fragments as a tool useful to identify and isolate structurally related polynucleotides. At low stringency hybridization conditions, for instance, polynucleotide libraries can be probed to identify genes that are at least about 50% homologous to the GLP-

2 receptor gene. To facilitate isolation of rat GLP-2 receptor gene homologs that are also GLP-2 receptor-encoding, stringency conditions are desirably enhanced to identify homologs having at least 80% (medium stringency) sequence identity homology at the polynucleotide level to receptor gene. More desirably the WBR gene homologs are 90% identical, (high stringency) and most desirably they have at least 95% sequence (high stringency) identity when compared to WBR. Preferably, the isolated WBR homologs are characterized in that (1) they can be amplified using the PCR primers of SEQ ID NO: 3 and SEQ ID NO: 4 and (2) they bind to the probe of SEQ ID NO: 5 under high stringency conditions.

Still more preferably, the isolated homologs are those which bind, under conditions of high stringency, with consensus regions of the GLP-2 receptor-encoding polynucleotides, i.e., those regions of the rat and human GLP-2 receptor-encoding polynucleotides which are identical and, relative to GLP-1 receptor-encoding sequences, are also unique to the GLP-2 receptor-encoding polynucleotides. Alignment of this nature reveals a number consensus regions for GLP-2: for example nucleotides spanning from 1460-1786. In one embodiment of the invention, these sequences, and their complements, constitute polynucleotide fragments useful, as just described for the intact gene, to identify polynucleotides structurally related to the human and rat GLP-2 receptor embodiments of the invention.

In a related embodiment, the cDNA sequence or the unique fragments of the GLP-2 receptor, most preferably of the human GLP-2 receptor, can be used in appropriately labelled form, e.g.,  $p^{32}$  labelled, for diagnosis of conditions associated with aberrant expression of the GLP-2 receptor. For example, over or under expression or expression in an inappropriate tissue. In one embodiment appropriate PCR primers (e.g., regions unique to the GLP-2 receptor but conserved between species) can be used diagnostically to identify aberrant structure or levels of GLP-2 receptor mRNA, e.g., associated with inherited or acquired disease states.

It has been found that the human GLP-2 receptor is located on chromosome 17P13. Thus, in a further embodiment, the invention provides expression products from this locus that

hybridize with the human GLP-2 receptor polynucleotide(Figure 6; SEQ ID NO:11) under stringent conditions.

As source material to isolate GLP-2 receptor-encoding homologs of the rat GLP-2 receptor-  
5 encoding polynucleotide gene, it is desirable but not necessary to use libraries of fetal or  
mature hypothalamal, jejunal, hindbrain or stomach tissue obtained from the vertebrate  
species targeted for receptor isolation. The invention accordingly includes not only the rat  
GLP-2 receptor-encoding polynucleotide of SEQ ID NO:1, but structural homologs thereof  
and particularly those that code for proteins having GLP-2 receptor properties. As  
10 exemplified hereinbelow, the WBR gene has been used successfully as a starting material  
to clone the human homolog of the rat GLP-2 receptor. Thus, the invention provides  
polynucleotides that encode GLP-2 receptors, including rat GLP-2 receptor and vertebrate  
homologs, particularly mammalian homologs thereof including human homologs, as well  
as synthetic variants of these.

15 It will be appreciated that such homologs can also be identified in libraries by screening, as  
noted, with fragments of the rat receptor gene or the human homolog, which incorporate at  
least 15 nucleotides, and preferably at least 25 nucleotides. With reference to SEQ ID NO: 1  
and the nucleotide numbering appearing thereon, suitable nucleotide fragments include in  
20 addition to the consensus nucleotide fragments noted above, those corresponding in sequence  
to the extracellular GLP-2 binding domain, and the stipulated transmembrane regions and the  
C-terminal portion of the receptor.

Technically, the identification of GLP-2 receptor genes can be achieved by applying standard  
25 hybridization or amplification techniques to a tissue-derived polynucleotide library. A wide  
variety of such libraries are commercially available. Where construction of a cDNA library is  
necessary, established techniques are applied. For example, isolation of such a WBR  
homolog typically will entail extraction of total messenger RNA from a fresh source of tissue,  
such as hypothalamal, jejunal, stomach or hindbrain tissue, preferably hypothalamal tissue or  
30 cell lines derived from these tissues, followed by conversion of message to cDNA and  
formation of a library in for example a bacterial plasmid, more typically a bacteriophage.

Such bacteriophage harboring fragments of the DNA are typically grown by plating on a lawn of susceptible *E. coli* bacteria, such that individual phage plaques or colonies can be isolated. The DNA carried by the phage colony is then typically immobilized on a nitro-cellulose or nylon-based hybridization membrane, and then hybridized, under carefully  
5 controlled conditions, to a radioactively (or otherwise) labelled probe sequence to identify the particular phage colony carrying the fragment of DNA of particular interest, in this case a rat or human GLP-2 homolog. The phage carrying the particular gene of interest is then purified away from all other phages from the library, in order that the foreign gene may be more easily characterized. Typically, the gene or a portion thereof is then isolated by subcloning  
10 into a plasmidic vector for convenience, especially with respect to the full determination of its DNA sequence.

As an alternative to obtaining GLP-2 encoding DNA directly as a DNA insert from an available or a constructed cDNA library, in light of the present disclosure it can be  
15 synthesized de novo using established techniques of gene synthesis. Because of the length of the GLP-2 receptor-encoding DNAs of SEQ ID NO: 1, SEQ ID NO: 9 and SEQ ID NO: 11, application of automated synthesis may require staged gene construction, in which regions of the gene up to about 300 nucleotides in length are synthesized individually and then ligated in correct succession for final assembly. Individually synthesized gene regions can be  
20 amplified by PCR. The application of automated synthesis may typically be applied by synthesizing specific regions or fragments of the gene and ligating them, usually via designed overlaps, in correct succession to form the final gene sequence. In this case, the longer the oligonucleotide building blocks, the fewer will be the ligations needed, resulting in greater ease of assembly.

25 The application of automated gene synthesis techniques provides an opportunity for generating sequence variants of the naturally occurring GLP-2 receptor gene. It will be appreciated, for example, that polynucleotides coding for the GLP-2 receptor herein described can be generated by substituting one or more synonymous codons for those represented in the  
30 naturally occurring polynucleotide sequences herein provided and such "synonymous codon equivalents" are within the scope of the present invention. In addition, polynucleotides

coding for synthetic variants of the GLP-2 receptor herein provided can be generated which incorporate from 1 to 20, e.g., from 1 to 5, amino acid substitutions, or deletions or additions. Preferred sites for such a modification include areas of non homology between the rat and human sequence, for example amino acid ranges 70-92, 328-350 and 475-504. Since it will be desirable typically to retain the natural ligand binding profile of the receptor for screening purposes, it is desirable to limit amino acid substitutions, for example to the so-called conservative replacements in which amino acids of similar charge are substituted (Fig. 9), and to limit substitutions to those sites less critical for receptor activity. For example, substitution of nucleotides "G" and "A" for nucleotides "A" and "G" respectively at positions 374 and 375 of the human cDNA sequence of SEQ ID NO: 11; resulting in the replacement of the naturally occurring arginine residue at position 85 of SEQ ID NO: 12 with a glutamic acid residue, provides a functional receptor. This functional receptor is referred to herein as the Glu<sup>85</sup> variant human GLP-2 receptor.

Having obtained GLP-2 receptor encoding polynucleotide, GLP-2 receptor can be produced in a number of ways, including in vitro transcription and via incorporation of the DNA into a suitable expression vector and expression in the appropriate host, for example in a bacterium such as E.coli, in yeast or in insect or in a mammalian cell. A variety of gene expression systems have been adapted for use with these hosts and are now commercially available, and any one of these systems can be selected to drive expression of the GLP-2 receptor-encoding DNA. Expression vectors may be selected to provide transformed cell lines that express the receptor-encoding DNA either transiently or in a stable manner. For transient expression, host cells are typically transformed with an expression vector harboring an origin of replication functional in a mammalian cell. For stable expression, such replication origins are unnecessary, but the vectors will typically harbor a gene coding for a product that confers on the transformants a survival advantage, to enable their selection such as a gene coding for neomycin resistance in which case the transformants are plated in medium supplemented with neomycin.

These expression systems, available typically but not exclusively in the form of plasmidic vectors, incorporate expression cassettes the functional components of which include DNA

constituting expression controlling sequences and optionally also signal peptides encoding sequences, which are host-recognized and enable expression of the receptor-encoding DNA when linked 5' thereof. The systems further incorporate DNA sequences which terminate expression when linked 3' of the receptor-encoding region. Thus, for expression in the  
5 selected mammalian cell host, there is generated a recombinant DNA expression construct in which the receptor-encoding DNA is linked with expression controlling DNA sequences recognized by the host, and which include a region 5' of the receptor-encoding DNA to drive expression, and a 3' region to terminate expression.

10 Included among the various recombinant DNA expression systems that can be used to achieve mammalian cell expression of the receptor-encoding DNA are those that exploit promoters of viruses that infect mammalian cells, such as the promoter from the cytomegalovirus (CMV), the Rous sarcoma virus (RSV), simian virus (SV40), murine mammary tumor virus (MMTV) and others. Also useful to drive expression are promoters  
15 such as the LTR of retroviruses, insect cell promoters such as those regulated by temperature, and isolated from *Drosophila*, as well as mammalian gene promoters such as those regulated by heavy metals i.e. the metallothionein gene promoter, and other steroid-inducible promoters.

20 In another of its aspects, the invention provides cells or membranes derived therefrom which are adapted by genetic alteration for use, for example, in identifying GLP-2 receptor ligands. In preferred embodiments, such cells are adapted genetically by the insertion of polynucleotide coding for a GLP-2 receptor. In particularly preferred  
25 embodiments, such cells incorporate a recombinant DNA molecule, e.g. an expression construct/vector, in which DNA coding for the GLP-2 receptor and expression controlling elements functional in the host are linked operably to drive expression of the DNA. For incorporation of receptor into cell plasma membranes, the vector can, if desired, be designed to provide a suitable heterologous signal peptide sequence to substitute for the signal peptide encoded naturally within the receptor DNA.

Suitable GLP-2 producing cells include the Chinese hamster ovary (CHO) cells for example of K1 lineage (ATCC CCL 61) including the Pro5 variant (ATCC CRL 1281); the fibroblast-like cells derived from SV40-transformed African Green monkey kidney of the CV-1 lineage (ATCC CCL 70), of the COS-1 lineage (ATCC CRL 1650) and of the COS-7 lineage (ATCC CRL 1651); murine L-cells, murine 3T3 cells (ATCC CRL 1658), murine C127 cells, human embryonic kidney cells of the 293 lineage (ATCC CRL 1573), human carcinoma cells including those of the HeLa lineage (ATCC CCL 2), and neuroblastoma cells of the lines IMR-32 (ATCC CCL 127), SK-N-MC (ATCC HTB 10) and SK-N-SH (ATCC HTB 11).

For use in ligand screening assays, cell lines expressing the receptor-encoding DNA can be stored frozen for later use. Such assays may be performed either with intact cells, or with membrane preparations derived from such cells. The membrane preparations typically provide a more convenient substrate for the ligand binding experiments, and are therefore preferred as binding substrates. To prepare membrane preparations for screening purpose, i.e., ligand binding experiments, frozen intact cells are homogenized while in cold water suspension and a membrane pellet is collected after centrifugation. The pellet is then washed in cold water, and dialyzed to remove any endogenous GLP-2 receptor ligands that would otherwise compete for binding in the assays. The dialyzed membranes may then be used as such, or after storage in lyophilized form, in the ligand binding assays.

The binding of a candidate ligand to a selected GLP-2 receptor of the invention can be assessed typically using a predetermined amount of cell-derived membrane (measured for example by protein determination), generally from about 25ug to 100ug. Generally, competitive binding assays will be useful to evaluate the affinity of a test compound relative to GLP-2. This competitive binding assay is performed by incubating the membrane preparation with radiolabelled GLP-2 peptide, for example [ $H^3$ ] or a radioiodinated GLP-2 analog, in the presence of unlabelled test compound added at varying concentrations. Following incubation, either displaced or bound radiolabelled GLP-2 can be recovered and measured, to determine the relative binding affinities of the test compound and GLP-2 for the



GLP-2 receptor used as substrate. In this way, the affinities of various compounds for the GLP-2 receptor can be measured.

Alternatively, binding of a candidate ligand to a GLP-2 receptor can be assessed using a functional assay. Using this approach, for example, intact cells harvested about two days following transient transfection or after about the same period following plating of stably transfected cells can be used to assess ligand binding. In a preferred embodiment, 293 EBNA cells (Invitrogen Cat. R620-07) are stably transformed with the pREP7 vector (Invitrogen Cat. V007-50) incorporating expressibly therein a GLP-2 receptor. Thereafter, binding of an agonist (or using a competition base format an antagonist) to the receptor can be discerned by measuring the level of intracellular cAMP. Most conveniently, intracellular cAMP is measured indirectly using a reporter system, wherein an easily measurable and preferably easily quantifiable downstream event indicates the level of intracellular cAMP. For example, measuring the level of the expression of a reporter gene construct having polynucleotide sequence under the control of a promoter which is responsive to cAMP. Alternatively, measurement of intracellular calcium, released from intracellular stores in response to an increase in intracellular cAMP, can be used as an indicator of the level of intracellular cAMP, for example, by incorporating into the transformed cell a protein that fluoresces on binding to calcium. In a preferred embodiment, intracellular cAMP levels are measured using the commercially available EIA kit. An additional advantage of the functionally based approach to assessing ligand binding is that the system can be automated allowing high throughput and ultra high through screening of vast chemical libraries.

As an alternative to using cells that express receptor-encoding DNA, ligand characterization may also be performed using cells for example *Xenopus* oocytes, that yield functional membrane-bound receptor following introduction of messenger RNA coding for the GLP-2 receptor. In this case, the GLP-2 receptor-encoding polynucleotide of the invention is typically subcloned into a plasmidic vector such that the introduced gene may be easily transcribed into RNA via an adjacent RNA transcription promoter supplied by the plasmidic vector, for example the T3 or T7 bacteriophage promoters. RNA is then transcribed from the inserted gene in vitro, and can then be injected into *Xenopus* oocytes. Each oocyte is a

single cell, but is large enough to be penetrated by a fine-tipped microneedle without causing irreparable damage. Following the injection of nL volumes of an RNA solution, the oocytes are left to incubate for up to several days, whereupon the oocytes are tested for the ability to respond to a particular ligand molecule supplied in a bathing solution.

Candidate GLP-2 receptor ligands can vary widely in structure, and most suitably include proteins which are highly related to GLP-2 itself in terms of amino acid sequence. For instance, the peptides disclosed in co-pending United States patent applications WO97/39031 and WO96/32414, incorporated herein by reference, may usefully be screened for GLP-2 receptor binding activity.

In addition to naturally occurring GLP-2 receptor sequences functional chimeric, receptors, incorporating portions of the GLP-2 receptor sequence and the polynucleotides encoding them are also embodiments of the invention. Functional chimeric GLP-2 receptors are constructed by combining the extracellular receptive sequences of a GLP-2 receptor with one or more of the transmembrane and intracellular segments of a known seven transmembrane G-protein coupled receptors for test purposes. This concept was demonstrated by Kobilka et al. (1988, Science 240:1310-1316) who created a series of chimeric  $\alpha 2$ - $\beta 2$  adrenergic receptors (AR) by inserting progressively greater amounts of  $\alpha 2$ -AR transmembrane sequence into  $\beta 2$ -AR. The binding activity of known agonists changed as the molecule shifted from having more  $\alpha 2$  than  $\beta 2$  conformation, and intermediate constructs demonstrated mixed specificity. The specificity for binding antagonists, however, correlated with the source of the transmembrane domain VII. The importance of transmembrane domain VII for ligand recognition was also found in chimeras utilizing two yeast  $\alpha$ -factor receptors and is significant because the yeast receptors are classified as miscellaneous receptors. Thus, functional role of specific domains appears to be preserved throughout the seven transmembrane G-protein coupled receptor family regardless of category.

In parallel fashion, internal segments or cytoplasmic domains from a particular GLP-2 receptor are exchanged with the analogous domains of a known seven transmembrane G-protein coupled receptor and used to identify the structural determinants responsible for coupling the

receptors to trimeric G-proteins (Dohlman et al. (1991) *Annu Rev Biochem* 60:653-688). A chimeric receptor in which domains V, VI, and the intracellular connecting loop from  $\beta 2$ -AR were substituted into  $\alpha 2$ -AR was shown to bind ligands with  $\alpha 2$ -AR specificity, but to stimulate adenylate cyclase in the manner of  $\beta 2$ -AR. This demonstrates that for adrenergic-type receptors, G-protein recognition is present in domains V and VI and their connecting loop. The opposite situation was predicted and observed for a chimera in which the V->VI loop from  $\alpha 1$ -AR replaced the corresponding domain on  $\beta 2$ -AR and the resulting receptor bound ligands with  $\beta 2$ -AR specificity and activated G-protein-mediated phosphatidylinositol turnover in the  $\alpha 1$ -AR manner. Finally, chimeras constructed from muscarinic receptors also demonstrated that V->VI loop is the major determinant for specificity of G-protein activity.

Chimeric or modified seven transmembrane G-protein coupled receptors containing substitutions in the extracellular and transmembrane regions have shown that these portions of the receptor determine ligand binding specificity. For example, two Ser residues conserved in domain V of all adrenergic and D catecholamine receptors are necessary for potent agonist activity. These serines are believed to form hydrogen bonds with the catechol moiety of the agonists within the binding site. Similarly, an Asp residue present in domain III of all seven transmembrane G-protein coupled receptors which bind biogenic amines is believed to form an ion pair with the ligand amine group in the binding site.

Functional, cloned seven transmembrane G-protein coupled receptors are expressed in heterologous expression systems and their biological activity assessed (e.g., Marullo et al. (1988) *Proc Natl Acad Sci* 85:7551-7555; King et al. (1990) *Science* 250:121-123). One heterologous system introduces genes for a mammalian seven transmembrane G-protein coupled receptors and a mammalian G-protein into yeast cells. The seven transmembrane G-protein coupled receptor is shown to have appropriate ligand specificity and affinity and trigger appropriate biological activation--growth arrest and morphological changes--of the yeast cells.

An alternate procedure for testing chimeric receptors is based on the procedure utilizing the  $P_{2u}$  purinergic receptor ( $P_{2u}$ ) as published by Erb et al. (1993, *Proc Natl Acad Sci* 90:10441-

104453). Function is easily tested in cultured K562 human leukaemia cells because these cells lack  $P_{2u}$  receptors. K562 cells are transfected with expression vectors containing either normal or chimeric  $P_{2u}$  and loaded with fura-a, fluorescent probe for  $Ca^{++}$ . Activation of properly assembled and functional  $P_{2u}$  receptors with extracellular UTP or ATP mobilizes intracellular  $Ca^{++}$  which reacts with fura-a and is measured spectrofluorimetrically. As with the seven transmembrane G-protein coupled receptors above, chimeric genes are created by combining sequences for extracellular receptive segments of any newly discovered seven transmembrane G-protein coupled receptors polypeptide with the nucleotides for the transmembrane and intracellular segments of the known  $P_{2u}$  molecule. Bathing the transfected K562 cells in microwells containing appropriate ligands triggers binding and fluorescent activity defining effectors of the seven transmembrane G-protein coupled receptors molecule. Once ligand and function are established, the  $P_{2u}$  system is useful for defining antagonists or inhibitors which block binding and prevent such fluorescent reactions.

In addition to using the receptor-encoding DNA to construct cell lines useful for ligand screening, expression of the DNA can according to another aspect of the invention be performed to produce fragments of the receptor in soluble form, for structure investigation, to raise antibodies and for other experimental uses. It is expected that the extracellular portion of the GLP-2 receptor contributes significantly to the binding of ligand molecule. It is therefore desirable in the first instance to facilitate the characterization of the receptor-ligand interaction by providing this extracellular ligand-binding domain in quantity and in isolated form, i.e., free from the remainder of the receptor. Such a construct has been made for the rat GLP-1 receptor, and it was shown to bind GLP-1 (Wilmen et al. (1996) FEBS LETTS, 398:43-47).

To accomplish this, the full-length GLP-2 receptor-encoding DNA may be modified by site-directed mutagenesis, so as to introduce a translational stop codon into the extracellular N-terminal region, immediately before the sequence encoding the first transmembrane domain (TM1), i.e., before residue 181 of SEQ ID NO: 2 and before residue 181 of SEQ ID NO: 12. Since there will no longer be produced any transmembrane domain(s) to "anchor" the receptor into the membrane, expression of the modified gene will result in the secretion, in

soluble form, of only the extracellular ligand-binding domain. Standard ligand-binding assays may then be performed to ascertain the degree of binding of a candidate compound to the extracellular domain so produced. It may of course be necessary, using site-directed mutagenesis, to produce several different versions of the extracellular regions, in order to optimize the degree of ligand binding to the isolated domains.

It will be appreciated that the production of such extracellular ligand binding domains may be accomplished in a variety of host cells. Mammalian cells such as CHO cells may be used for this purpose, the expression typically being driven by an expression promoter capable of high-level expression, for example the CMV (cytomegalovirus) promoter. Alternately, non-mammalian cells, such as insect Sf 9 (*Spodoptera frugiperda*) cells may be used, with the expression typically being driven by expression promoters of the baculovirus, for example the strong, late polyhedron protein promoter. Filamentous fungal expression systems may also be used to secrete large quantities of such extracellular domains of the GLP-2 receptor.

*Aspergillus nidulans*, for example, with the expression being driven by the *alcA* promoter, would constitute such an acceptable system. In addition to such expression hosts, it will be further appreciated that any prokaryotic or other eukaryotic expression system capable of expressing heterologous genes or gene fragments, whether intracellularly or extracellularly would be similarly acceptable.

The availability of isolated extracellular ligand-binding domains of the receptor protein makes it feasible to determine the 3-dimensional structures of these ligand-binding regions, with or without a candidate ligand complexed thereto, by a combination of X-ray crystallographic and advanced 2D-NMR techniques. In this way, additional new candidate compounds, predicted to have the required interactions with the 3-dimensional receptor structure, can be specifically designed and tested.

With large domains, crystallography is the method of choice for structure determination of both the domain in isolation, and of the co-complex with the natural ligand (or an appropriate antagonist or agonist molecule). If a particular domain can be made small enough, for example approximately 100-130 amino acids in length, then the powerful technique of 2-D

NMR can also be applied to structure determination. This enables not only the determination of the domain structure, but also provides dynamic information about the drug-receptor interaction.

5 For use particularly in detecting the presence and/ or location, for example in intestinal tissue, the present invention also provides, in another of its aspects, labelled antibody to a GLP-2 receptor. To raise such antibodies, there may be used as immunogen either the intact, soluble receptor or an immunogenic fragment thereof, produced in a microbial or mammalian cell host as described above or by standard peptide synthesis techniques. Regions of the GLP-2  
10 receptor particularly suitable for use as immunogenic fragments include those corresponding in sequence to an extracellular region of the receptor, or a portion of the extracellular region, such as peptides consisting of 10 or more amino acids of the 401-509 region of SEQ ID NO: 2. With regard to the human GLP-2 receptor (SEQ ID NO: 12), peptides comprising the mature extracellular domain (residues 65-180); intracellular loop 3 (residues 363-385) and the  
15 intracellular C-terminal domain (residues 442-533) may be usefully employed as immunogens for the production of antibodies to the human GLP-2 receptor.

Antibodies to the desired GLP-2 receptor or fragment immunogen are available, for polyclonal antibody production, from the blood of an animal that has been immunized with  
20 the immunogen. Alternatively, for monoclonal antibody production, immunocytes such as splenocytes can be recovered from the immunized animal and fused, using hybridoma technology, to a myeloma cells. The fusion products are then screened by culturing in a selection medium, and cells producing antibody are recovered for continuous growth, and antibody recovery. Recovered antibody can then be coupled covalently to a detectable label,  
25 such as a radiolabel, enzyme label, luminescent label or the like, using linker technology established for this purpose.

Animal model systems which elucidate the physiological and behavioral roles of the GLP-2 receptor are produced by creating transgenic animals in which the activity of the  
30 GLP-2 receptor is either increased or decreased, or the amino acid sequence of the expressed GLP-2 receptor is altered, by a variety of techniques. Examples of these

techniques include, but are not limited to: 1) Insertion of normal or mutant versions of DNA encoding a GLP-2 receptor, by microinjection, electroporation, retroviral transfection or other means well known to those skilled in the art, into appropriate fertilized embryos in order to produce a transgenic animal or 2) Homologous recombination of mutant or normal, human or animal versions of these genes with the native gene locus in transgenic animals to alter the regulation of expression or the structure of these GLP-2 receptor sequences. The technique of homologous recombination is well known in the art. It replaces the native gene with the inserted gene and so is useful for producing an animal that cannot express native GLP-2 receptors but does express, for example, an inserted mutant GLP-2 receptor, which has replaced the native GLP-2 receptor in the animal's genome by recombination, resulting in under expression of the transporter. Microinjection adds genes to the genome, but does not remove them, and so is useful for producing an animal which expresses its own and added GLP-2 receptors, resulting in over expression of the GLP-2 receptors.

One means available for producing a transgenic animal, with a mouse as an example, is as follows: Female mice are mated, and the resulting fertilized eggs are dissected out of their oviducts. The eggs are stored in an appropriate medium such as M2 medium. DNA or cDNA encoding a GLP-2 receptor is cesium chloride purified from a vector by methods well known in the art. Inducible promoters may be fused with the coding region of the DNA to provide an experimental means to regulate expression of the transgene. Alternatively or in addition, tissue specific regulatory elements may be fused with the coding region to permit tissue-specific expression of the trans-gene. The DNA, in an appropriately buffered solution, is put into a microinjection needle (which may be made from capillary tubing using a piper puller) and the egg to be injected is put in a depression slide. The needle is inserted into the pronucleus of the egg, and the DNA solution is injected. The injected egg is then transferred into the oviduct of a pseudopregnant mouse ( a mouse stimulated by the appropriate hormones to maintain pregnancy but which is not actually pregnant), where it proceeds to the uterus, implants, and develops to term. As noted above, microinjection is not the only methods for inserting DNA into the egg cell, and is used here only for exemplary purposes.

The invention having been described above, may be better understood by referring to the following examples. The following examples are offered for the purpose of illustrating the invention and should not be interpreted as a limitation of the invention.

## EXAMPLE 1

### Isolation of the GLP-2 receptor

#### PCR-assisted cloning of partial rat and mouse GLP-2 receptor cDNAs

Rat Neonate Intestine cDNA library (Stratagene, La Jolla, CA; Cat. 936508) and Mouse Jejunum first strand cDNA was prepared. Degenerate primers M-2F/S (SEQ ID NO: 3) and M-7R/S (SEQ ID NO: 4) were used to amplify a partial fragment of the rat GLP-2 receptor from the Rat Neonate Intestine cDNA library and of the mouse GLP-2 receptor from Mouse Jejunum template. The protocol is described below:

#### Degenerate PCR :

6 µl 10x VENT buffer from New England Biolabs

6 µl 2.5µM each stock dATP, dCTP, dGTP and dTTP

4 µl rat neonate intestine cDNA (1:10 dilution)

3 µl 25µM M2F/S primer [5'-TTTTCTAGAASRTSATSTACACNGTSGGCTAC-3']  
(SEQ ID NO: 3)

3 µl 25µM M7R/S primer [5'-TTTTCTCGAGCCARCARCCASSWRTARTTGGC-3']  
(SEQ ID NO: 4)

2 µl (10 units) Amplitaq DNA polymerase (Perkin Elmer)

36 µl ddH<sub>2</sub>O.

Reaction conditions: 35 cycles at 94°C, 2°min.; 94°C, 1 min.; 53°C, 30 sec.; 72°C, 1 min.

The predominant PCR product was a 303 base pair (bp) DNA fragment. 30 µl samples of the above PCR were purified using the QIAGEN PCR purification kit and eluted in 30 µl ddH<sub>2</sub>O. The resulting product was then re-amplified using the same degenerate PCR conditions, with the exception only of 31 cycles at 94°C.



The predominant product at 303 base pair (bp) was cut out and purified using QIAGEN QIAquick gel purification protocol into 30 µl ddH<sub>2</sub>O. The resulting product was then re-amplified using the same degenerate PCR conditions, with the exception only of 31 cycles at 94°C.

5

Next, double digest (Xba I and Xho I) was done on the entire reamplified PCR reaction as follows: 28 µl DNA; 16 µl 10X One-Phor-All buffer (Pharmacia); 2 µl (40 units) Xba I enzyme (Pharmacia); 2 µl (40 units) Xho I enzyme (Pharmacia); and 30 µl ddH<sub>2</sub>O.

10 The samples were digested 4 hours in 37°C water block heater, brought up to 100 µl volume with ddH<sub>2</sub>O (sterile) and purified by (1) equal amount (100ul) chloroform extraction; (2) weekend precipitation with 2 volumes ethanol/10 volumes 3M sodium acetate; (3) 1x wash with 70% EtOH; and (4) resuspension in 10 µl 1x TE (pH 8.0).

15 pBluescript clone 5HT1F#9 was next digested with Xba I and Xho I as follows:

10 µl DNA (pBluescript clone 5HT1F#9)

5 µl 10X NEBuffer 2 (New England Biolabs)

3 µl (1:20 dilution=3 units) Xba I (New England Biolabs)

3 µl (1:20 dilution=3 units) Xho I (New England Biolabs)

20 5 µl (10x) BSA (New England Biolabs)

24 µl ddH<sub>2</sub>O.

25 The sample was digested for 3 hours in 37 C water block heater, heat-inactivated at 65°C for 20 min and purified using GeneCleanII kit from BIO 101. Aliquots of the PCR reactions were cloned into the above pBluescript plasmid vector using T4 DNA ligase kit (New England Biolabs) and transformed into Epicurean Coli XL-2 Blue MRF' Ultracompetent cells (Stratagene). The transformation was plated onto 2xYT + AMP plates and single colonies were picked. DNA minipreps were made using QIAGEN QIA-prep 8 miniprep kit and the sequences of the fragments were determined using ABI  
30 system. Novel sequences were identified containing a partial fragment of the rat and mouse GLP-2 receptor sequence.

Cloning of cDNA with complete GLP-2 receptor coding region was achieved as follows:  
First, cDNA libraries from the following three tissues were used for screening,

1. Rat Hypothalamus (RHT)
2. Rat Hind Brain (RHB)
- 5 3. Rat Duodenum and Jejunum (RDJ)

The three cDNA libraries were prepared by priming with random primer and subcloning unidirectionally into Hind III and Not I sites of pcDNA3.

10 Next, the three cDNA libraries were homology screened by a degenerate oligo C4-4 [5'-AACTACATCCACMKGMAYCTGTTYVYGTCBTTCATSCT-3'] (SEQ ID NO: 5) by colony lifts and filter hybridization. The following hybridization conditions were employed: 5X SSPE (1X SSPE is 0.18M NaCl, 10mM NaH<sub>2</sub>PO<sub>4</sub> (pH 7.4), 10mM EDTA (pH 7.4)) and 5X Denharts solution (1% Ficoll, 1% Polyvinylpyrrolidone, 1% BSA); 25 mg/ml salmon sperm DNA.

15 Filters were hybridized at 50°C overnight. Then the filters were washed 2 times in 2X SSPE and 1% SDS at room temperature for 30 min, 2 times in 2X SSPE and 1% SDS at 50°C for 20 min per wash, and finally two times in 1X SSPE and 0.5% SDS. Positive clones were identified by autoradiography. A plug of 1cm<sup>2</sup> surrounding the positive  
20 clone was removed from the plate and placed in 1 ml of 2x YT +20% Glycerol, vortexed and was frozen at -80°C.

25 Plasmid DNA from positive plugs was prepared as follows: 100 ml of bacterial culture of each positive plug was grown on an agar plate. The bacterial cells were scraped and resuspended in 1 ml of 2xYT medium+20% Glycerol. Bacterial pellet from the 250 ml of bacterial resuspension was resuspended in 150 ml of Solution I (50mM Glucose, 10mM Tris-HCl, 1mM EDTA), lyse in Solution II (0.2M NaOH, 1% SDS), neutralized with ice cold Solution III (Potassium acetate; 4 vol. of 5M potassium acetate + 1 vol. of 10M acetic acid). After pelleting bacterial DNA, 340 ml isopropanol was added to the  
30 supernatant. This was centrifuged at max for 15 min. The pellet was resuspended in TE + 20 mg/ml RNase, incubated at 37 C for 30 minutes and precipitated with isopropanol

+ 0.2M potassium acetate. After centrifugation, the pellet was washed with 70% alcohol, allowed to air dry and resuspended in TE.

Plasmid DNA from 2777-clone pools of rat hypothalamus cDNA library RHT cDNA library was next exploited as follows: Two primers were designed from an area of the PCR-cloned GLP-2 receptor cDNA sequence that did not have identity to known receptors of the gene family. The two primers P23-R1 and P23-F1 amplified a 196 bp fragment only from novel clone DNA but not with GLP-1 receptor cDNA or PACAP receptor cDNA. The Expand<sup>TM</sup> PCR system from Boehringer Mannheim (Cat. 1681-842) was used under the following conditions:

2 µl of 10x Expand<sup>TM</sup> Buffer 1

2.8 µl of 2.5mM dNTP mix

0.6 µl of primer P23-R1 [5'-TCATCTCCCTCTTCTTGGCTCTTAC-3'] (SEQ ID NO: 6)

0.6 µl of primer P23-F1 [5'-TCTGACAGATATGACATCCATCCAC-3'] (SEQ ID NO: 7)

0.3 µl of Expand PCR enzyme (1 unit)

12.7 µl water

1 µl DNA

Reaction conditions: 32 cycles at 93°C, 40 sec; (cycles) 58°C, 40 sec; (cycles) 68°C, 40 sec

DNAs from each positive plug or pool of 2777-clone pools were amplified with specific primers P23-F1 and P23-R1 under the conditions specified above. Out of 1057 C4-4 hybridization-positive plugs and 884 2777-clone pools only five template sources amplified a 196 bp PCR product. These were: (1) Plug 334, (2) Plug 780, (3) RHT pool 233, (4) RHT pool 440, and (5) RHT pool 587.

Amplification of GLP-2R cDNA from the five positive templates was then performed. By using one specific primer (P23-R1 or P23-F1) and one primer based on pcDNA3 vector (Invitrogen) sequence (830F or 1186R), the GLP-2R cDNA insert was directly

amplified from clonally impure plugs or 2777-clone pools. The sequences of the vector primers were as follows:

830F: [5'-AACCCACTGCTTAC-3']

1186R: [5'-CCCAGAATAGAATGACACC-3']

5 The PCR was done, under the following conditions just noted, using Expand™ PCR system.

10 The most prominent band was reamplified, purified and sequenced. Based on the amplified sequence obtained, additional primers were designed and new sequencing carried out. In this manner the complete sequences of the GLP-2R cDNA inserts in all five sources of clones were determined. Sequence analysis showed that only pool RHT 440 and pool RHT 587 contain clones with complete coding sequence of GLP-2R and that the two clones were identical (derived from the same cDNA clone).

15 Because of difficulty in clonally purifying the GLP-2 receptor cDNA clone from the RHT 440 or RHT 587 cDNA library pools, the cDNA was amplified and recloned into pcDNA3. Based on the sequence obtained from RHT 440 and RHT 587, two primers were designed one which primed starting 4bp upstream of the initiation codon and another which primed starting 8bp downstream of the stop codon.

20 WBR-C5: [5'-CAGGGGCCGGTACCTCTCCACTCC-3']

WBR-C3: [5'-TTGGGTCCTCGAGTGGCCAAGCTGCG-3']

25 The two primers were used to amplify a DNA fragment of approximately 1525 bp fragment under the following PCR conditions using Expand™ PCR system from Boehringer Mannheim (Catalogue no. 1681-842).

10 µl of 10x Expand™ PCR Buffer 1

14 µl of 2.5mM dNTP mix

3.0 µl of Primer 1 (10 µM) (WBR-C5)

30 3.0 µl of Primer 2 (10 µM) (WBR-C3)

1.5 µl of Enzyme (5 units)

63.5 µl water

5 µl DNA

Reaction conditions: 5 cycles ( 93°C, 1 Min; 72°C, 40s; 60°C, 45 sec; 68°C, 2 min) and 25 cycles ( 93°C, 1 min; 72°C, 1 min; 68°C, 2 min).

The amplified product was subcloned into Kpn I and Xho I sites of pcDNA3 vector (Invitrogen). Plasmid DNA was prepared using the method described above.

## EXAMPLE 2

### Functional Assay

Cos-1 cells were transfected as described in Analytical Biochemistry, 218:460-463(1994) with Rat clone 587 GLP-2 receptor, cloned human GLP-2 receptor (pC3/HuGL2R-2), or cloned residue 85 variant human GLP-2 receptor (pC3.1/HuGL2R-MH4), pcDNA3. Rat GLP-1 (7-36) amide was used as a control peptide. Solutions used were as follows:  
RSC in RPMI 1640 (49ml RPMI + 1ml FCS + 50ul chloroquine, 100mM);  
DEAE/RSC Solution: 18.4 ml RSC +1.6 ml DEAE/Dextran (10 mg/ml).

The assay procedure entailed the following:

a) 50 mg of either rat clone 587 GLP-2 receptor, or cloned human GLP-2 receptor, or cloned residue 85 variant human GLP-2 receptor was added (as plasmid pcDNA3) to a 50 ml tube containing six mls. of RSC and incubated at 37°C.

b) Six ml of DEAE/RSC solution was added to each tube and incubated at 37°C for 2 min.

c) 1.5 ml of COS-1 cell suspension (5.5 millions cells) was added to each tube and incubated for 1 hr 45 min at 37°C.

d) Following incubation, the sample was spun for 5 minutes at low speed, washed with DMEM/F12 + 10% FBS twice, and the pellet resuspended in 12.5 ml DMEM/F12 + 10% FBS media.

e) One ml of cell suspension (step d) was added to each well of 6 well plates coated with poly- D-lysine (from Collaborative Biomedical), containing 3 ml of media (0.45 million cells/well).

f) Plates were incubated at 37°C for 3 days.

Treatment of Transfected Cos-1 cells with GLP-1/GLP-2 analog was done as follows:

Solutions : DMEM/F12 (SFM)+ IBMX (3-isobutyl-1-methylxanthin) 0.85mM +0.1% ascorbic acid and 10 um pargyline (all solutions purchased from Sigma). Media was prepared fresh on day of use.

Assay Procedure: The culture media of each well (transfected 6 well plates, cells) was removed, and the wells were washed once with SFM media. Then 2 ml of SFM + IBMX media was added to each well and plates were incubated at 37°C for 10 min.

Following incubation, the SFM + IBMX was removed from each well and fresh SFM + IBMX media containing GLP-1/GLP-2 (GLP-1,7-36,amide from Sigma, [Gly2]hGLP-2 from Allelix) concentration 1, 3, 10 and 30 nM were added to the appropriate wells.

Plates incubated at 37°C incubator for 30 min. Following incubation, the media were removed from each well. The wells were washed once with 1ml PBS (Phosphate Buffered Saline). Each well was then treated with 1ml cold 95% ethanol: 5mM EDTA (2:1) at 4°C for 1 hr. Cells from each well then were scraped and transferred into individual eppendorf tubes. Tubes were centrifuged for 5 min at 4°C, and the supernatants were transferred to new eppendorf tubes and dried in speed vacuum.

Following drying, tubes were reconstituted in 100ul of Na- Acetate and kept at 4°C, 25 µl of this solution used for cAMP Assay.

The functional assay was performed as follows: cAMP content for each extract was determined in duplicate by EIA (Enzyme ImmunoAssay) using the Amersham Biotrak cAMP EIA Kit (Amersham 225). Results of the assays, illustrated in Figure 3 and Figure 8, demonstrate the GLP-2 selectivity exhibited by the cloned rat and human receptors. In a similar functional assay used to users binding to the GLP-1 receptor, the expected specificity for GLP-1 was observed.

### EXAMPLE 3

#### Isolation Of Human GLP-2 Receptor cDNA

#### Medium-stringency hybridization screening of a human hypothalamus cDNA library

One million clones from a  $\lambda$ gt10 cDNA library from human hypothalamus (Clontech; Cat. No. 1172a) were screened by plaque lifts on nitrocellulose filters (Amersham; Cat.RPN137E). The probe was prepared by random primer labelling of a DNA fragment containing the complete coding region of rat GLP-2 receptor. The DNA fragment was isolated from clone 587-C1, which contains the complete coding region from SEQ ID NO: 2.

Pre-hybridization and hybridization were each carried out overnight in a hybridization solution consisting of 50% formamide, 5X SSPE, 5X Denhart's solution, 0.5% SDS and salmon sperm DNA (200 mg/ml). After hybridization the filters were washed under the following conditions (medium stringency):

two times at room temperature in 2X SSPE and 0.01% SDS.

two times at 42°C in 2X SSPE and 0.01% SDS.

two times at 42°C in 0.2X SSPE and 0.01% SDS.

The filters were autoradiographed and agar plugs, each containing numerous bacteriophage plaques, were picked from regions on the plates corresponding to positive signals on the filter. From one million cDNA clones sampled in the first round screen, there were identified two positive clones (HHP6-1 and HHP13). On secondary screening only HHP13 turned out positive. Several positive plaques (HHS13) from the HHP13 plate were pooled and taken for tertiary screening. Three single positive plaques from this round of screening were picked (HHT13-1, HHT13-2, HHT13-3).

PCR amplification was then used for partial sequencing of the positive clones. On a lawn of bacterial cells (E.coli C600Hfl), 10  $\mu$ l of phage suspension from each clone was applied at marked spots. After 5 hr incubation at 37°C, the phage plaques were clearly visible and covered  $\sim 1$  cm<sup>2</sup>. A portion of each plaque was transferred to 200  $\mu$ l of water. The samples were incubated in a boiling water bath for 5 min and centrifuged at room temperature for 10 min. One millilitre of sample was used for PCR amplification with two sets of degenerate primers:

M2FS[5'-TTTTTCTAGAASRTSATSTACACNGTSGGCTAC-3'] (SEQ ID NO: 3) and

M7RS[5'-TTTTCTCGAGCCARCARCCASSWRTARTTGGC-3'] (SEQ ID NO: 4); or  
C4-4 [5'-AACTACATCCACMKGMAYCTGTTYVYGTCBTTCATSCT-3'] (SEQ ID  
NO: 5) and

C9-2R [5'-TCYRNCTGSACCTCMYYRTTGASRAARCAGTA-3'] (SEQ ID NO: 8).

The Expand™ PCR system from Boehringer Mannheim (Cat. 1681-842) was used under  
the following conditions:

5 µl of 10x Expand™ Buffer 3

7 µl of 2.5mM dNTP mix

1.5 µl of primer M2FS or C4-4

1.5 µl of primer M7RS (with M2FS) or C9-2R (with C4-4)

0.75 µl of Expand PCR enzyme (1 unit)

33.25 µl water and

1 µl DNA.

Reaction conditions were: 32 cycles at 93°C, 1 min; cycles 50 °C, 1 min; cycles 45 °C, 1  
min; cycles 68°C, 2 min.

M2F/S and M7R/S amplified a DNA fragment of about 300 bp and C4-4 and C92-R  
amplified a DNA fragment of about 700 bp. The PCR products were purified using the  
QIAGEN QIAquick PCR purification kit (Cat. 28104) and eluted in 50 µl 10 mM Tris,  
pH 8.0. Sequence analysis of the products revealed no differences between the  
templates, as expected from the fact that they represent multiple copies of a single cDNA  
clone (HHT13).

A number of factors indicate that this clone contains coding sequence of the human  
GLP-2 receptor. One factor is the degree of sequence similarity. The glucagon receptor  
cDNA can be used to predict the expected degree of sequence conservation found  
between rat and human receptors. At the nucleotide level, there is 82.6% identity within  
the coding regions of the rat and human glucagon receptors. At the amino acid level,  
there is 80.9% identity and 89.1% amino acid similarity between the glucagon receptors  
of the two species.



In the case of the human GLP-2 receptor cloned herein, the sequence of the partial human GLP-2 receptor cDNA (HHT13) is highly homologous to rat GLP-2 receptor cDNA at both the nucleotide and amino acid level. SEQ ID NO: 9 shows 87.1% identity with the rat GLP-2 receptor cDNA sequence. The predicted amino acid sequence of this cDNA region has 87.4% identity and 93.2% similarity with the predicted amino acid sequence of the rat GLP-2 receptor. The total predicted length of the rat receptor preprotein is 550 amino acids, suggesting about 44% of the coding region of the human receptor had been identified.

Further evidence supporting this conclusion comes from a comparison of the partial human GLP-2 receptor amino acid sequence with the rat GLP-2 receptor and the 3 next closest family members, shown below:

Receptor Sequence (amino acid)	Percent Identity with HHT13	Percent Similarity
GLP-2 receptor (rat)	87.4	93.2
GLP-1 receptor (rat)	50.0	74.1
Glucagon receptor (rat)	51.4	73.9
GIP receptor (rat)	50.7	70.3

These comparisons, together with the benchmark provided by sequence similarities between the rat and human glucagon receptors, provide definitive evidence that the cDNA designated HHT13 represents a fragment of the human counterpart of the rat GLP-2 receptor.

The full amino acid sequence of the human GLP-2 receptor can be obtained by first determining the sequence of the complete cDNA inserts in HHT13-1, HHT13-2 and HHT13-3. By using degenerate primers for PCR amplification and subsequent sequencing, we obtained sequence from only part of each insert. It is possible that these identical clones contain an insert which spans the complete coding sequence of the human GLP-2 receptor preprotein. To determine the complete sequence of the cDNA insert, the clones are grown in large quantity to prepare approximately 20 mg of each

equivalent clone. The complete cDNA insert is excised by restriction with Eco RI, and subcloned into pcDNA3 (Invitrogen). Alternatively, two primers from vector sequence flanking the insert are used to amplify the complete cDNA insert using the Expand<sup>TM</sup> PCR system from Boehringer Mannheim (Cat. 1681-842). The amplified cDNA is cut with appropriate restriction enzymes and is subcloned into pcDNA3 (Invitrogen).

If a complete coding sequence is not present in the HHT13 clones, cDNA libraries are screened for additional clones to complete the coding region of human GLP-2 receptor cDNA. Preferably human cDNA libraries (from Stratagene or Clontech) representing the following tissues are used for screening: Human hypothalamus; Human fetal brain; Human duodenum and jejunum; Human stomach; and Human fetal intestine.

Two PCR primers are designed from the sequence of human GLP-2 receptor cDNA already determined. These primers are designed such that they could not amplify any related gene family members other than the GLP-2 receptor cDNA itself. A dilution of the cDNA library stock is used to make library sub-pools such that 50,000 clones are represented in each pool. PCR is conducted with the GLP-2 receptor-specific primers to diagnose pools containing a GLP-2 receptor cDNA clone, using the Expand<sup>TM</sup> PCR system from Boehringer Mannheim (Cat. 1681-842) under the following conditions:

2 µl of 10x Expand<sup>TM</sup> Buffer 1

2.8 µl of 2.5mM dNTP mix

0.6 µl of primer P1

0.6 µl of primer P2

0.3 µl of Expand PCR enzyme (1 unit)

12.7 µl water

1 µl of library pool containing 50,000 clones

Reaction conditions: 32 cycles at 93°C, 40 sec; 50-58°C, 40 sec; 68°C, 40 sec.

Sequence is then obtained from the complete GLP-2 receptor cDNA insert from a positive pool. By using one specific primer and one primer based on vector sequence close to the cloning site, the GLP-2 receptor cDNA insert is directly amplified from

clonally impure clone pools, using the Expand™ PCR system from Boehringer Mannheim (Catalogue no. 1681-842) most suitably under the following conditions:

2 µl of 10x Expand™ Buffer 1

2.8 µl of 2.5mM dNTP mix

5 0.6 µl of Primer 1

0.6 µl of Primer 2

0.3 µl of Enzyme (1 unit)

12.7 µl water

1 µl of library pool stock

10 Reaction conditions: 32 cycles at 93°C, 45 sec; 50°C, 45 sec; 68°C, 1 min.

The reaction is run on a preparative agarose gel, and the most prominent band is purified and sequenced. Based on the amplified sequence obtained, additional primers are designed to obtain sequence and clones of complete coding region and clone the complete cDNA

15

5' RACE and 3' RACE are used to obtain complete coding sequence of the human GLP-2 receptor cDNA. Rapid Amplification of cDNA Ends (RACE) is a procedure routinely used for amplification of DNA sequences from first cDNA strand (easily prepared from mRNA) template between a defined internal site and either 3' or the 5' end of the mRNA. Total or mRNA from different human tissues are commercially available from Clontech. The 3' RACE System (Gibco-BRL Life Technologies; Cat. 18373-019) and 5' RACE System (Cat. 18374-058) kits are used. The manuals of these two products provide detailed protocols. In brief, protocols are as described below.

20

25

For the 3' RACE procedure, first strand cDNA synthesis is initiated at the poly (A) tail of mRNA using the adapter primer (provided with system) incorporating a unique sequence for universal PCR amplification of the RACE products. After synthesis of the first strand cDNA from this primer, the original mRNA template is destroyed with RNase H. Amplification is then performed using two primers: one is a gene-specific primer (which will be designed from the available partial cDNA sequence of HHT13);

30

the other is the universal amplification primer provided with the kit. The amplified product is subcloned into a plasmid vector for sequencing.

For the 5' RACE System, the first strand cDNA is synthesized from mRNA using a gene-specific primer (which is based on the available partial cDNA sequence of HHT13) and SuperScript II reverse transcriptase. The original mRNA template is removed by treatment with RNase H. Unincorporated dNTPs, primer, and proteins are separated from cDNA using spin cartridge. A homopolymeric dCTP tail is then added to the 3'-end of the first strand cDNA using TdT enzyme and dCTP nucleotides. PCR amplification is performed using two primers: one is a nested, gene-specific primer designed from the available partial DNA sequence of HHT13; and the other is an "anchor primer" provided with the system. Both primers incorporate restriction sites for subcloning into plasmids and subsequent sequencing.

Sub-cloning of HHT13  $\lambda$ gt10 clones into pcDNA3, their sequencing and expression

A. Amplification of cDNA inserts with  $\lambda$ gt10 primers.

On a lawn of bacterial cells (E.coli C600Hfl), 10  $\mu$ l of phage resuspension from each clone was placed at marked spots. After 5 hr incubation at 37°C, the phage plaques were clearly visible. The surface of each plaque was transferred to 200  $\mu$ l of water. The samples were kept in boiling water bath for 5 minutes and centrifuged at room temperature for 10 minutes. 1  $\mu$ l of sample was used to amplify with a set of  $\lambda$ gt10 primers.

GT10-5KXb [5'-GGGTAGTCGGTACCTCTAGAGCAAGTTCAGCC-3']

vs

GT10-3BXh [5'-ATAACAGAGGATCCTCGAGTATTTCTTCCAG-3']

The Expand™ PCR system from Boehringer Mannheim (Cat. 1681-842) was used under the following conditions:

5  $\mu$ l of 10x Expand™ Buffer 3

7  $\mu$ l of 2.5mM dNTP mix

1.5 µl of primer GT10-5KXb  
1.5 µl of primer GT10-3BXh  
0.75 µl of Expand PCR enzyme (1 unit)  
33.25 µl water  
5 1 µl DNA

Reaction conditions: 5 cycles of 93°C-40 sec; 50 °C- 1 min; 68°C-2 min and 30 cycles of 93°C-40 sec; Ramp to 68°C- 1 min; 68°C-2 min.

10 An amplified DNA fragment of about 2200 bp long was seen on the agarose gel from all three clones. The PCR product were purified using the QIAGEN's QIAquick PCR purification kit (Cat. no. 28104) and eluted in 50 µl 10 mM Tris, pH 8.0. The templates were sequenced.

#### B. Subcloning into pcDNA3 vector.

15 The amplified and purified DNA from the three clones was restricted with Kpn I and Xho I and subcloned into pcDNA3 restricted with similar restriction enzymes. The plasmids were named pHHT13-1, pHHT13-2, and pHHT13-3. Plasmids DNAs were prepared using either crude method (alkaline treatment, bacterial DNA precipitation with 3 M KOAc, isopropanol precipitation followed by RNase treatment and second round of  
20 isopropanol precipitation) or plasmid DNA kits from Qiagen Inc. The templates prepared using Qiagen's kits were sequenced.

#### C. Functional Assay

25 Transfections were carried out with each clone, using the rat GLP-2R, 587 clone as a positive control for cAMP response to GLP-2 peptide. Methods for transfection, cell culture and cAMP assay were identical to those described for the functional assay of rat, 587 clone. Results showed that although the positive control gave good cAMP response in COS cells, none of the HHT13 clones gave any cAMP response. As confirmed by sequencing which showed a frame-shift mutation, the functional data suggested that no  
30 functional GLP-2R protein was expressed from these cDNA clones.

D. Comparison of DNA sequences between rat GLP-2R and HHT13 subclones.

The comparison showed a 2 bp deletion at a position corresponding to nucleotides 389-390 of the rat GLP-2R cDNA, resulting in the loss of nucleotides 374-375 of the human GLP-2R cDNA sequence presented herein.

PCR was used to incorporate two bp of the rat GLP-2R DNA sequence into HHT13-1 DNA at the site of the 2 bp frame-shift deletion identified relative to the rat GLP-2R coding sequence. The following primers were designed from HHT13 DNA sequence to insert two bp :

HWBR/2BPI-475F

[5'-ACAGGCATGTCTGGAAGACTTACTCAAGGAACCTTCTGGCAT-3']

HWBR/2BPI-506R

[5'-ATGCCAGAAGGTTCCCTTGAGTAAGTCTTCCAGACATGCCTGT-3']

HWBR-F7 [5'-TTCCTCTGTGGTACCAAGAGGAATGC-3']

and HWBR-1910R:

[5'-GGTGGACTCGAGGTACCGATCTCACTCTCTTCCAGAATC-3']

PCR 1: One ng of pHHT13-1 DNA was used as template to do two PCRs with primers, HWBR-F7 vs HWBR/2BPI-506R and HWBR/2BPI-475F vs HWBR-1910R.

The Expand™ PCR system from Boehringer Mannheim (Cat. 1681-842) was used under the following conditions:

5 µl of 10x Expand™ Buffer 1

7 µl of 2.5mM dNTP mix

1.5 µl of primer HWBR-F7 or HWBR/2BPI-475F

1.5 µl of primer HWBR/2BPI-506R or HWBR-1910R

0.75 µl of Expand PCR enzyme (1 unit)

33.25 µl water and

1 µl DNA.

Reaction conditions: 10 cycles of 92°C-40 sec; 48°C- 1 min; 68°C-3 min and 30 cycles of 92°C-40 sec; 55°C- 40 sec; 68°C-2 min.

The primers HWBR-F7 and HWBR/2BPI-506R amplified a DNA fragment of 400 bp and HWBR/2BPI-475F and HWBR-1910R amplified a DNA fragment of approximately 1.4 kb on an agarose gel. The two bands were cut out of the agarose gel and purified with Qiaquick gel extraction kit from Qiagen Inc. (Cat no. 28706) and the DNAs were eluted in 50 µl of 10 mM Tris (pH 8.5).

PCR 2 (Extension without primers): Approximately 75 ng of two amplified product from above PCR 1 were mixed and then recombined without primers by extending under the following conditions:

2 µl of 10x Expand™ Buffer 1  
2.8 µl of 2.5mM dNTP mix  
0.3 µl of Expand PCR enzyme (1 unit)  
8.9 µl water  
6 µl of combined PCR 1 products

Reaction conditions: 15 cycles of 92°C-1 min; 60°C- 5 min; 68°C-3 min.

PCR 3: 1 µl of amplified mix from PCR 2 was used as template to amplify with HWBR-F7 and HWBR- 1910R primers using the following conditions:

10 µl of 10x Expand™ Buffer 1  
14 µl of 2.5mM dNTP mix  
3.0 µl of primer HWBR-F7 or HWBR/2BPI-475F  
3.0 µl of primer HWBR/2BPI-506R or HWBR-1910R  
1.5 µl of Expand PCR enzyme (1 unit)  
67.5 µl water and  
1 µl DNA.

Reaction conditions: 30 cycles of 92°C-1 min; 60°C- 1 min; 68°C-2 min.

A DNA fragment of approximately 1.7 kb was amplified as seen on an agarose gel. The PCR product was purified using the QIAGEN's QIAquick PCR purification kit (cat. no. 28104) and eluted in 50 µl of 10 mM Tris, pH 8.0. The purified product was restricted with Kpn I and subcloned into Kpn I-restricted pcDNA3.1(-)/Myc-His A (Invitrogen,

Cat. no. V855-20). One clone, named pc3.1/HuGL2R/MH6 (pHuMH6), had the 1.7 kb insert in correct orientation as checked by PCR using vector vs. insert primers.

Functional assay:

This hybrid clone was compared to rat GLP-2R using the assay described in Example 2. Results showed that the 2 bp "GA" replacement into the putative deletion site yielded a clone encoding a functional GLP-2R protein, as shown by the cAMP response to GLP-2 treatment.

**EXAMPLE 6**

**Isolation of the Full-Length Human GLP-2 Receptor cDNA**

Twenty thousand clones from  $\lambda$ gt10 cDNA Library from Human Stomach (Clontech; Cat. HL3017a) were plated on each of 100 agar 150 mm plates. SM buffer (0.1 M NaCl, 10 mM  $Mg_2SO_4$ , 35 mM Tris, pH-7.5, 0.01% gelatin) was added to each plate to obtain 100 phage lysates each containing 20,000 (20K) pooled clones. The first fifty 20K phage lysates (20K pools) were screened by PCR using two primers designed from HHT13 DNA sequence. The template DNA from each pool was prepared by boiling phage lysate for 10 minutes and centrifuging for 10 minutes.

HWBR-113F [5'-GTGGAGAGGATTTGTGCAAACATTTC-3']

HWBR-578R [5'-AGAGACATTTCAGGAGAAGAATGAG-3']

1  $\mu$ l of each 20K pool DNA was diagnosed by PCR with HWBR-113F and HWBR-578R primers using the following conditions:

2  $\mu$ l of 10x Expand<sup>TM</sup> Buffer 1

2.8  $\mu$ l of 2.5mM dNTP mix

0.6  $\mu$ l of primer HWBR-113F

0.6  $\mu$ l of primer HWBR-578R

0.3  $\mu$ l of Expand PCR enzyme (1 unit)

12.7  $\mu$ l water

1  $\mu$ l 20K pool DNA

Reaction conditions: 35 cycles of 92°C- 40s.; 60°C- 40s.; 68°C-1 min.



A DNA fragment of approximately 450 bp was seen in amplification of templates from two pools (HST 19 and HST 38).

B. Screening of clones from two positive pools: HST 19 and HST 38.

40,000 clones plated from each of two positive 20K pools were screened by plaque lifts on nitrocellulose filters (Amersham; Cat.RPN137E). The probe was prepared by random primer labelling a DNA fragment from pHHT13-1.

1. The filters were pre-hybridized and hybridized at 42°C overnight. Hybridization solution consisted of 50% formamide, 5X SSPE, 5X Denhart's solution, 0.5% SDS and salmon sperm DNA (200 mg/ml).

2. After hybridization the filters were washed under the following conditions:

- two times at room temperature in 2X SSPE and 0.01% SDS;
- two times at 42°C in 2X SSPE and 0.01% SDS; and
- two times at 50°C in 0.1X SSPE and 0.01% SDS.

3. The filters were autoradiographed and the regions on the plates matching to positive signals were isolated. One positive clone (HST 38-4-30) was isolated from HST 38 pool. 450 bp DNA fragment was amplified from the positive clone by using primers HWBR-113F and HWBR-578R and sequenced. The sequence clearly showed that the plasmid contain 2 bp (AG) at position 373-374 of HHT13 DNA sequence.

The complete insert of clone HST 38-4-30 was amplified using  $\lambda$ gt10 primers as described in Example 1. PCR amplified a DNA fragment of approximately 1.4 kb. The amplified DNA was purified and sequenced.

#### **EXAMPLE 7**

##### **Reconstruction of a clone of full-length functional human GLP-2R cDNA and functional assay**

A 700 bp fragment obtained by Kpn I and Pvu II restriction digest of the amplified DNA from clone HST 38-4-30, and 1.4 kb DNA fragment from Xho I and Pvu II restricted pHHT13-1 DNA were subcloned into Kpn I and Xho I restricted pcDNA3 in a three-way ligation. The new plasmid construct was called pc3/HuGL2R-2. In this manner the full length sequence of the human GLP-2 receptor was obtained.

#### Functional Assay:

The new clone was compared to the rat GLP-2R clone 587 as described previously above. Results showed that the clone encoded a functional human GLP-2R protein, which led to cAMP production in COS cells in response to GLP-2 treatment (Figure 8).

### **EXAMPLE 8 Antibodies directed to the GLP-2 receptor**

#### **1. Antipeptide antibodies**

Antipeptide antibodies were raised in rabbits against an N-terminal peptide (QTRENTTDIWQDESE), a C-terminal peptide (SEG DGSETLQKLR) and extracellular loop 1 (SHNSUSKRPDDESG) of the rat GLP-2 receptor.

Immunocytochemical analysis of serum produced as above confirmed that serum contained antibodies directed to the GLP-2 receptor which do not cross react with the rat GLP-1 receptor.

#### **2. Antibodies to a GLP-2 receptor raised against a fusion protein**

Polynucleotide encoding the C-terminal region of rat GLP-2 receptor (amino acids 444-550) was spliced to the C-terminus of glutathione S-transferase (GST) in pGEX-2T and expressed in E.coli strain SUREI. Protein was purified using affinity chromatography using the above GLP-2 C-terminal fragment fused to the C-terminus of maltose binding protein. Protease degradation was minimized by using a cocktail of protease inhibitors (Boeringer Mannheim).

Antibodies were raised generally according to the method disclosed in Antibodies: A laboratory manual, Harlow and Lane, Cold Spring Harbor Laboratory, 1988. Briefly, the

5 GLP-2-GST fusion protein was used to raise antibodies in rabbits as follows. Initial injection was with 100 µg of fusion protein in complete Freund's adjuvant at multiple sites, intramuscularly and subcutaneously. Booster injections were made at multiple sites intramuscularly with 100 µg of fusion protein in incomplete Freund's adjuvant at days 14, 21, 42 and 56.

10 Antisera was affinity purified using a GLP-2-MBP fusion protein affinity column. Immunocytochemical analysis confirmed that these antibodies specifically recognize the GLP-2 receptor.

### EQUIVALENTS

15 The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. Indeed, various modifications of the above-described makes for carrying out the invention which are obvious to those skilled in the field of biochemistry, molecular biology or related fields are intended to be within the scope of the following claims.

What is claimed is:

1. An isolated polynucleotide encoding a GLP-2 receptor.
- 5 2. The isolated polynucleotide according to claim 1, which codes for the human GLP-2 receptor having the amino acid sequence of amino acids 67-553 of SEQ ID NO:12
3. The isolated polynucleotide according to claim 2, comprising nucleotides 318-1780 of  
10 SEQ ID NO:11
4. The isolated polynucleotide according to claim 1, which codes for the rat GLP-2 receptor having the amino acid sequence of amino acid 67-550 of SEQ ID NO:2
5. The isolated polynucleotide according to claim 4, having nucleotides 335-1789 of  
15 SEQ ID NO: 1.
6. An isolated polynucleotide which hybridizes under conditions of high stringency with a polynucleotide defined in claim 3 or claim 5.
- 20 7. An oligonucleotide comprising at least 15 nucleotides and having a polynucleotide sequence that hybridizes selectively to a polynucleotide according to any of claims 1-6.
8. An oligonucleotide according to claim 7, wherein the nucleotide sequence of the oligonucleotide corresponds identically with a region of the polynucleotide defined in  
25 claim 3 or claim 5.
9. An oligonucleotide according to claim 7, wherein the nucleotide sequence of the oligonucleotide corresponds identically with a region common to the polynucleotides defined in claim 3 and claim 5.
- 30

10. In labelled form, a polynucleotide selected from a polynucleotide as defined in any of claims 1-6, and an oligonucleotide as defined in any of claims 7-9.

5 11. A recombinant polynucleotide comprising a GLP-2 receptor-encoding polynucleotide as defined in any of claims 1- 6, and expression controlling elements linked operably therewith to drive expression thereof.

12. A cell that has been genetically engineered by the incorporation expressibly therein of a polynucleotide according to any of claims 1-6.

10 13. The cell according to claim 11, which is a mammalian cell.

14. An antibody which selectively binds to a GLP-2 receptor.

15 15. A recombinant GLP-2 receptor.

16. The GLP-2 receptor of claim 15, comprising amino acids 67-553 of SEQ ID NO:2.

17. The GLP-2 receptor of claim 15, comprising amino acids 67-550 of SEQ ID NO: 12

20 18. A method for identifying GLP-2 receptor ligands, which comprises the steps of:

25 (1) incubating a candidate ligand with a cell as defined in claim 12 or with a membrane preparation obtained therefrom; and then

(2) determining whether binding between the GLP 2 receptor and the candidate ligand has occurred.

30 19. A GLP-2 receptor ligand identified by the method of claim 18.

1/12

1 AAGCTTCGCG GCCTCTGCAG AKGACTTGTG CAAACACTTC CTCTCTGGAC  
51 AAGGAGGAAT GCAGGAGGCC ACCGCCTGCA GTACATCTTG GAGTGTGGGA  
101 GGGATGTGCC TGCACCTGTG AACGGGCGCC AGGAGAATGA GGGCCCAACC  
151 AAGCCCGGCA GTGCCCAGTA GATGCAGAGA GGCACCCGTG CCCCAGTGA  
201 GGGCACAGCC AGTGGGCATC CCTGAGGCCC AGGGGCCCGT TCCTCTCCAC  
251 TCCCAACAGA TGCGTCTGCT GTGGGGCCCT GGGAGGCCCT TCCTCGCCCT  
301 GCTTCTGCTG GTTTCATCA AGCAAGTTAC AGGATCGCTC CTCAAGGAGA  
351 CAACTCAGAA GTGGGCTAAT TATAAGGAGA AGTGTCTGGA AGACTTGCAC  
401 AATAGACTTT CTGGCATATT TTGTAATGGG ACATTTGATC GGTATGTGTG  
451 CTGGCCTCAT TCTTATCCTG GAAATGTCTC TGTTCCCTGT CCTTCATACT  
501 TACCTTGGTG GAATGCAGAG AGCCCAGGAA GGGCCTACAG AACTTGCTTG  
551 GCTCAGGGGA CTTGGCAGAC GCGAGAGAAC ACCACAGATA TTTGGCAGGA  
601 TGAATCAGAA TGCTCAGAGA ACCACAGCTT CAGACAAAAC GTGGATCACT  
651 ACGCCTTGCT ATACACCTTG CAGCTGATGT AACTGTGGG CTACTCCGTG  
701 TCTCTCATCT CCTCTTCTT GGCTCTTACA CTCTTCTTGT TCCTTCGAAA  
751 ACTGCATTGC ACACGCAATT ACATCCACAT GAACCTGTTT GCTTCGTTCA  
801 TCCTGAAAGT TCTGGCTGTC CTGGTGAAGG ACATGGTCTC CCACAACCTT  
851 TACTCCAAGA GGGCCGATGA TGAGAGTGGA TGGATGTCAT ATCTGTCAGA  
901 GACATCCGTC TCCTGTGCTT CCGTCCAGGT CCTCCTGCAC TACTTTGTGG  
951 GCACCAATCA CTTGTGGCTG CTGGTTGAAG GACTTTACCT CCACACTCTG  
1001 CTGGAGCCCA CAGTGTTCCT TGAAAGGCGG CTGTGGCCCA AGTACCTGGT  
1051 GGTGGGTTGG GCCTTCCCA TGCTGTTTGT TATTCCTGG GGTTCGCCC  
1101 GTGCACACCT GGAGAACACA CGGTGCTGGG CCACAAATGG GAACCTGAAA  
1151 ATCTGGTGGA TCATCAGAGG ACCCATGCTG CTTGTGTAA CAGTTAATT  
1201 CTTCACTTTC CTCAAGATTC TCAAGCTTCT CATTTCTAAG CTCAAAGCTC  
1251 ATCAGATGTG CTTCAGAGAC TACAAATACA GATTGGCGAA ATCAACGTTG  
1301 CTCCTCATTC CTTGTGTTGG GGTTCATGAG GTCCTCTTCA CTTTCTTCCC  
1351 CGACGACCAA GTTCAAGGAT TTTCAAACG TATTCGACTC TTCATCCAGC  
1401 TGACACTGAG CTCTGTCCAC GGATTTCTGG TGGCCTTGCA GTATGGCTTT

1451 GCCAATGGAG AGGTGAAGGC AGAGCTGCGA AAGTCATGGG GCCGCTTCTT  
1501 ATTAGCCCGC CACTGGGGCT GCAGAACCTG TGTCCTGGGG AAGAATTTCC  
1551 GGTTCCTGGG GAAGTGTTC AAGAAGCTGT CGGAGGGAGA TGGCTCTGAG  
1601 AACTCCAGA AGCTGCGGTT CTCCACATGC AGCTCACACC TGGCCTCTGA  
1651 GACCCTGGGA GACGTTGGGG TACAGCCTCA CAGGGGCCGT GGAGCTTGGC  
1701 CCCGGGGAAG CAGCCTGTCT GAGAGCAGTG AGGGAGACTT CACCCTGGCC  
1751 AATACGATGG AGGAGATTCT GGAAGAGAGT GAGATCTAAG GCAGGGTCCA  
1801 TCACCGCAGC TTGGCCACTG ARGAmCCAAC CcTARGAAGG ATKTTGCCGA  
1851 RCCCARGGTC CTCCTCTTCC TATGTWcAT MCCCATTTTG ATGTGAAGTC  
1901 TCTCCTGGGT GAMCAASCTC TGTACCAACS ARTCTCagTC CCTCTTGCCC  
1951 TTGTCAcCCT ACTACCCCTC CCCCATCaCa CATgTTTTCC AGAATKTCGG  
2001 TTGGTTTGGG GGGGGGGGTC TTGCCCTAAA TTCAAGTsGA GTGGARCCCA  
2051 CCATGAAGAA AARTCATTTA TTAAATAGAR TCCGGTTAGG ATCTCCTTCC  
2101 CGTTCATGGT GCATGGCCTC CTTCCAAGGG ATGGGAGTCG GSTGCACTGG  
2151 AACCCACAG GAAAcTTTGA AGTATCCAGT TCTAGGGAAT TATAGCCAAT  
2201 ATTCTGAGAG AGCAAGTCTG AGATGAGAKC CGAGAATWGC AAGTGTWGA  
2251 CAWGCATTCA AGGAAACTCC TCACCTTTgG GCGAAACCTA tGGCAGGATc  
2301 GGCAcGGAGC AGCTATTMTG CAAYGGCCGC TCACCTGGGA CATACCACTC  
2351 TCCTTGGGCA GGATGTGACC CCATGTKGTC CCCCAGACTC CTCTCCTCCT  
2401 TGCTTSTSTT CYTTTCYGT CAAGTCTCAC CTCCCTTCT ACATCTCAGT  
2451 TCWGTTTGGT GTYGACAGAA GYYTGAATGT CACAATACTG CATGTGTTAG  
2501 TTTCTGTCGT CATTGCTGTG TCCAAATACC TGACCAGGAC CAATTTAAGC  
2551 GAGGAACTGC TACATGGGCG GCCGC


FIGURE 1

1 MRPQSPPAVP SRCREAPVPR VRAQFVGIP E AQGFVPLHSQ QMRLLWGPGR  
51 PFLALLLLVS IKQVTGSLK ETTQKWANYK EKCLEDLHNR LSGIFCNGTF  
101 DRYVCWPHSY PGNVSVPCPS YLPWWNAESP GRAYRHCLAQ GTWQTRENTT  
151 DIWQDESECS ENHSFRQNV D HYALLYTLQL MYTVGYSVSL ISLFLALTLF  
201 LFLRKLHCTR NYIHMNLFAS FILKVLAVLV KDMVSHNSYS KRPDDESGWM  
251 SYLSETSVSC RSVQVLLHYF VGTNHLWLLV EGLYLHTLLE PTVFPERRLW  
301 PKYLVVGWAF PMLFVIPWGF ARAHLENTRC WATNGNLKIW WIIRGPMLLC  
351 VTVNFFIFLK ILKLLISKLK AHQMCFRDYK YRLAKSTLLL IPLLGVEVL  
401 FTFPPDDQVQ GFSKRIRLFI QLTLLSSVHGF LVALQYGFAN GEVKAELRKS  
451 WGRFLLARHW GCRTCVLGKN FRFLGKCSKK LSEG DGSETL QKLRFSTCSS  
501 HLASETLGDV GVQPHRGRGA WPRGSSLSES SEGDFTLANT MEEILEEESI

FIGURE 2



Figure 3

cAMP Response of COS Transfected with Clone

HGLP2 Human GLP-2 Receptor C4-4 vs C9-2R PCR from Clone HHT13

```
1  TCGTTCTCTC TTATCTCCCT GTTCTGGGT CTCACCTCC TCTTCTTCT
51  TCGAAACTC CACTGCAGGC GCAACTACAT CCACATGAAC TTGTTTGCTT
101 CTTTCATCCT GAGAACCCTG GCTGTACTGG TGAAGGACGT CGTCTTCTAC
151 AACTCTTACT CCAAGAGGCC TGACAATGAG AATGGGTGGA TGTCTTACCT
201 GTCAGAGATG TCCACCTCCT GCCGCTCAGT CCAGGTTCTC TTGCATTACT
251 TTGTGGGTGC CAATTACTTA TGGCTGCTGG TTGAAGGCTT CTACCTCCAC
301 ACGCTGCTGG AGCCACACAGT GCTTCCTGAG AGGCGGCTGT GGGCCARATA
351 CCTGCTGTTG GCTTGGGCTT TCCCTGTGCT ATTTGTTGTA CCCTGGGGTT
401 TCGCCCGTGC ACACCTGGAR AACACAGGCT GCTGGACAA ACATGGGGAT
451 AAGAAATCT GGTGGATCAT CCGAGGACCC ATGATGCTCT GTGTAACAT
501 CAATTTCTTC ATCTTCCTGA AATTTCTCAA GCTTCTCATT TCTAAGCTCA
551 AAGCTCATCA AATGTGCTTC AGAGATTATA AATACAGATT GGCAAAATCA
601 ACACTGGTCC TCATTCCTTT ATTGGGCTTT CATGAGATCC TCTTCTCTTT
651 CATCACTGAT GATCAAG
```

Figure 4

6/12

1	S F S L I S L F L A L T L L L F L R K L TCCTTCTCTCTTATCTCCCTCTTCCTGGCTCTGACCTCTCTTCTTTCTTCGAAACTC -----+	50
61	H C T R N Y I H M N L F A S F I L R T L CACTGCACGGCGCAACTACATCCACATGAACCTTGTGCTTCTTTCATCTCTGAGAACCTG -----+	120
121	A V L V K D V V F Y N S Y S K R P D N E GCTGTACTGGTGAAGGACGTCGTCTTCTACAACTCTTACTCCAAGAGGCTGACAAATGAG -----+	180
181	N G W M S Y L S E M S T S C R S V Q V L AATGGGTGGATGTCTACCTGTGAGAGATGTCCACCTCTGCGGCTCACTCCAGGTTCTC -----+	240
241	L H Y F V G A N Y L W L L V E G L Y L H TTGCATTACTTTGTGGGTGCCAATTACTTATGGTCTGCTGTTGAAGGCTCTACCTCCAC -----+	300
301	T L L E P T V L P E R R L W P K Y L L L ACGCTGCTGGAGCCCGACATGCTTCTGAGAGGCGGCTGTGCGCCCATACCTGCTGTTG -----+	360
361	G W A F P V L F V V P W G F A R A H L E GTTTGGGCTTCCCTGTGCTATTTGTTGTACCTGGGCTTTCGCGCTGCACACCTGGAR -----+	420
421	N T G C W T T N G N K X I W H I I R G P AACACAGGGTGCTGGACACAAATGGGAATAAGAAAATCTGTGGATCATCCGAGGAGCC -----+	480
481	M H L C V T V N F F I F L H I L N L L I ATGATGCTCTGTGTACACATCAATTTCTTCATCTTCTGAAAATCTCAAGCTTCTCAT -----+	540
541	S K L K A H Q M C F R D Y K Y R L A K S TCTAAGCTCAAAGCTCATCAATGTGCTTCAGAGATTATAAATACAGATTGGCAAAATCA -----+	600
501	T L V L I P L L G V H E I L F S F I T D ACACTGGTCTCTATTCTTTATTGGGCTTTCATGAGATCTCTCTCTTTCATCACTGAT -----+	660
661	D Q GATCAAG ----- 557	

Figure 5

7/12

(SEQ ID NO: 11)

1 TGGAGAGGATTGTGCAAACATTTCTTCTGTGGACCAAGAGGAATGCAAGAGGAGGCTGC 50  
-----+-----  
61 CTGCGGTGCATCTTGGACGGCTAGAGAGATGTACCCCTACTTGTGAAGGTGCACGAGGAA 120  
-----+-----  
121 M K L G S S R A G P G R G S A G L L P G 180  
GATGAAGCTGGGATCGAGCAGGGCAGGGCCTGGGAGAGGAAGCGCGGGACTCCTGCCTGG  
-----+-----  
181 V H E L P M G I P A P W G T S P L S F H 240  
CGTCCACGAGCTGCCCATGGGCATCCCTGCCCTGGGGGACCAGTCCTCTCTCCTTCCA  
-----+-----  
241 R K C S L W A P G R P F L T L V L L V S 300  
CAGGAAGTGCTCTCTCTGGGCCCCCTGGGAGGCCCTTCCTCACTCTGGTCTCTGCTGGTTTC  
-----+-----  
301 I K Q V T G S L L E E T T R K W A Q Y K 360  
CATCAAGCAAGTTACAGGATCCCTCCTTGAGGAACGACTCGGAAGTGGGCTCAGTACAA  
-----+-----  
361 Q A C L R D L L K E P S G I F C N G T F 420  
ACAGGCATGTCTGAGAGACTTACTCAAGGAACCTTCTGGCATATTTTGTAAACGGGACATT  
-----+-----  
421 D Q Y V C W P H S S P G N V S V P C P S 480  
TGATCAGTACGTGTGTGGCCTCATCTCTCTGGAAATGTCTCTGTACCCTGCCCTTC  
-----+-----  
481 Y L P W W S E E S S G R A Y R H C L A Q 540  
ATACCTTACCTTGGTGGAGTGAAGAGAGCTCAGGAAGGGCCTACAGACACTGCTTGGCTCA  
-----+-----  
541 G T W Q T I E N A T D I W Q D D S E C S 600  
GGGGACTTGGCAGACGATAGAGAACGCCACGGATATTTGGCAGGATGACTCCGAATGCTC  
-----+-----  
601 E N H S F K Q N V D R Y A L L S T L Q L 660  
CGAGAACCACAGCTTCAAGCAAAACGTGGACCTTATGCCTTGCTGTCAACCTTGACGCT  
-----+-----  
661 M Y T V G Y S F S L I S L F L A L T L L 720  
GATGTACACCGTGGGATACTCCTTCTCTTATCTCCCTCTTCTGGCTCTCACCCCTCT  
-----+-----  
721 L F L R K L H C T R N Y I H M N L F A S 780  
CTTGTTTCTTCGAAAACCTCCACTGCACGGCAACTACATCCACATGAACCTGTTTGCTTC  
-----+-----  
781 F I L R T L A V L V K D V V F Y N S Y S 840  
TTTCATCCTGAGAACCCTGGCTGTACTGGTGAAGGACGTGCTCTTCTACAACCTTACTC  
-----+-----  
841 K R P D N E N G W M S Y L S E M S T S C 900  
CAAGAGGCCTGACAATGAGAATGGGTGGATGTCCTACCTGTCAGAGATGTCCACCTCCTG  
-----+-----  
901 R S V Q V L L H Y ~~E~~ V G A N Y L W L L V 960  
CCGCTCAGTCCAGGTTCTCTTGCACTTACTTATGGGTGCCAATTACTTATGGCTGCTGGT  
-----+-----  
961 E G L Y L H T L L E P T V L P E R R L W 1020  
TGAAGGCCTCTACCTCCACACGCTGCTGGAGCCACAGTGCTTCTGAGAGGCGGCTGTG  
-----+-----

FIGURE 6A

	P R Y L L L G W A F P V L F V V P W G F	
1021	GCCCAGATACCTGCTGTGGTGGGTCCTCGTGCTATTGTGTACCCTGGGGTTT -----+-----+-----+-----+-----+	1030
	A R A H L E N T G C W T T N G N K K I W	
1091	CGCCCGTGCAACCTGGAGAACAACAGGGTGTGGACAACAATGGGAATAAGAAAATCTG -----+-----+-----+-----+-----+	1140
	W I I R G P M M L C V T V N F F I F L K	
1141	GTGGATCATCCGAGGACCCATGATGCTCTGTGTAACAGTCAATTTCTTCATCTTCCTGAA -----+-----+-----+-----+-----+	1200
	I L K L L I S K L K A H Q M C F R D Y K	
1201	AATTCTCAAGCTTCTCATTCTAAGCTCAAAGCTCATCAAATGTGCTTCAGAGATTATAA -----+-----+-----+-----+-----+	1260
	Y R L A K S T L V L I P L L G V H E I L	
1261	ATACAGATTGGCAAATCAACTGGTCTCTCATTCTTTATTGGGCGTTTCATGAGATCCT -----+-----+-----+-----+-----+	1320
	F S F I T D D Q V E G F A K L I R L F I	
1321	CTTCTCTTTTCATCACTGATGATCAAGTTGAAGGATTGCAAAACTTATACGACTTTTCAT -----+-----+-----+-----+-----+	1380
	Q L T L S S F H G F L V A L Q Y G F A N	
1381	TCAGTTGACACTGAGCTCCTTTTCATGGGTTCTGGTGGCCTTGCGATATGGTTTTGCCAA -----+-----+-----+-----+-----+	1440
	G E V K A E L R K Y W V R F L L A R H S	
1441	TGGAGAAGTGAAGGCTGAGCTGCGGAAATACTGGGTCCGCTTCTTGCTAGCCCGCCACTC -----+-----+-----+-----+-----+	1500
	G C R A C V L G K D F R F L G K C P K K	
1501	AGGCTGCAGAGCCTGTGTCTGGGGAAGGACTTCCGGTTCCTAGGAAAATGTCCCAAGAA -----+-----+-----+-----+-----+	1560
	L S E G D G A E K L R K L Q P S L N S G	
1561	GCTCTCGGAAGGAGATGGCGCTGAGAAGCTTCGGAAGCTGCAGCCCTCACTTAACAGTGG -----+-----+-----+-----+-----+	1620
	R L L H L A M R G L G E L G A Q P Q Q D	
1621	GCGGCTCCTACATCTAGCCATGCGAGGTCTTGGGGAGCTGGGCGCCAGCCCCAACAGGA -----+-----+-----+-----+-----+	1680
	H A R W P R G S S L S E C S E G D V T M	
1681	CCATGCACGCTGGCCCCGGGGCAGCAGCCTGTCCGAGTGCAGTGAGGGGGATGTCAACCAT -----+-----+-----+-----+-----+	1740
	A N T M E E I L E E S E I *	
1741	GGCCAACACCATGGAGGAGATTCTGGAAGAGAGTGAGATCTAGGGTGGAGTTCACCACC -----+-----+-----+-----+-----+	1800
	CTGGCTCTGCTCCCAGGGA CTCTTGAGGGGGCCCGGAAGAGGAAGCAAAGCAGGACACA	
1801	-----+-----+-----+-----+-----+	1860
	CGTTGCTGGGCACGGAATCATTCTCGTTCCATTACCATGCCACTTTGATATGAAAGCTA	
1861	-----+-----+-----+-----+-----+	1920
	TCACAAGGTTCTTCAAGCTCTGTATGAAGAGGCTGTGTGTCATGCTCACAGCCTCTGCC	
1921	-----+-----+-----+-----+-----+	1980
	TGCTCTTCTCATCCTAATAACCCCCACCACTGTGTTTTCCACAATGCCCACCAGACCCTA	
1981	-----+-----+-----+-----+-----+	2040
	GGGCCTGGCTCTAAATTCAAGCCAATGAAGTCCACCCCGGAATTCTTTTGCTTTTACCC	
2041	-----+-----+-----+-----+-----+	2100
	CTGGAAGAAATA	
2101	-----+-----+-----+-----+-----+	2112

FIGURE 6B

9/12

Human GLP-2 Receptor Complete Open Reading Frame  
Note: Translation may start with M-1 or M-26.  
Length: 553 April 21, 1997 07:42 Type: P Check: 2775

```
1  MXLGSSRAGP GRGSAGLLPG VHELPMGIPA PWGTSPLSFH RKCSLWAFGR
51  PFLTLVLLVS IKQVTGSILLE ETTRKWAQYK QACLRDLLKE PSGIFCNGTF
101 DQYVCWPHSS PGNVSVPCPS YLPWWSEESS GRAYRHCLAQ GTWQTIENTAT
151 DIWQDDSECS ENHSFKQNV DRYALLSTLQL MYTVGYSFSL ISLFLALTLL
201 LFLRXLHCTR NYIHMLNFAS FILRTLAVLV KDVVVFYNSYS KRPDNENGWM
251 SYLSEMSTSC RSVQVLLHYF VGANYLWLLV EGLYLHTLLE PTVLPERRLW
301 PRYLLLGWAF PVLFFVPWGF ARAHLENTGC WTTNGNKKIW WIIRGPMMLC
351 VTVNFFIFLK ILKLLISKLK AHQMCFRDYK YRLAKSTLVL IPLLGVHEIL
401 FSFITDDQVE GFAXLIRLFI QLTLSSTHGF LVALQYGFAN GEVKAELRKY
451 WVRFLLARHS GCRACVLGKD FRFLGKCPKK LSEGDAEKL RKLQPSLNSG
501 RLLHLAMRGL GELGAQPQOD HARWPRGSSL SECSEGDVTM ANTMEEILEE
551 SEI
```

FIGURE 7

cAMP Response to GLP-2




FIGURE 8

11/12

```

1 MRPQSPPAVPSRCREAPVPRVRAQPVGIPEAQGPVPLHSQQMRLLWGPGR 50
|: | | | | . | | | | | | | | | | | | | |
1 MKLGSSRAGPGRGSAGLLPGVHELPMGIPAPWGTSPLSFHRKCSLWAPGR 50

51 PFLALLLLVSIKQVTGSLLKETTQKWANYKEKCLEDLHNRLSGIFCNGTF 100
| | | | . | | | | | | | | | | | | | | | | | | | | | | |
51 PFLTLLVLLVSIKQVTGSLLLEETTRKWAQYKQACLRDLLKEPSGIFCNGTF 100

101 DRYVCWPHSYPGNVSVPCPSYLPWWNAESPGRAYRHCLAQGTWQTRENTT 150
|. | | | | | | | | | | | | | | | | | | | | | | | |
101 DQYVCWPHSSPGNVSVPCPSYLPWWSEESSGRAYRHCLAQGTWQTIENTAT 150

151 DIWQDESECSSENHSFRQNVDPHYALLYTLQLMYTVGYSVSLISLFLALTTF 200
| | | | : | | | | | | | | | | | | | | | | | | | | | |
151 DIWQDDSECSSENHSFKQNVDRYALLSTLQLMYTVGYSFSLISLFLALTLL 200

201 LFLRKLHCTRNYIHMNLFASFILKVLAVLVKDMVSHNSYSKRPDDESGWM 250
| | | | | | | | | | | | | | | | : | | | | | | | | | | | |
201 LFLRKLHCTRNYIHMNLFASFILRTLAVLVKDVFYNSYSKRPDNENGWM 250

251 SYLSETSVSCRSVQVLLHYFVGTNHLWLLVEGLYLHTLLEPTVFPERRLW 300
| | | | | | | | | | | | | | | | : | | | | | | | | | | | |
251 SYLSEMSTSCRSVQVLLHYFVGANYLWLLVEGLYLHTLLEPTVLPERRLW 300

301 PKYLVVGWAFFPMLFVIPWGFARAHLENTRCWATNGNLKIWWIIRGPMMLC 350
| : | | . | | | | . | | | : | | | | | | | | | | | | | | : | |
301 PRYLLLGWAFFPVLFPVWGFARAHLENTGCWTTNGNKKIWWIIRGPMMLC 350

351 VTVNFFIFLKILKLLISKLKAHQMCFRDYKYRLAKSTLLLIPLLGVHEVL 400
| | | | | | | | | | | | | | | | | | | | | | | | | | | | : |
351 VTVNFFIFLKILKLLISKLKAHQMCFRDYKYRLAKSTLVLIPLLGVHEIL 400

401 FTFFPDDQVQGFSKRIRLFIQTLTSSVHGFLVALQYGFANGEVKAELRKS 450
|. | | | | : | | . | | | | | | | | | | | | | | | | | | |
401 FSFITDDQVEGFAKLIRLFIQTLTSSFHGFLVALQYGFANGEVKAELRKY 450

451 WGRFLLARHWGCRTCVLGKNFRFLGKCSKKLSEGDGSETLQKLRFSTCSS 500
| | | | | | | | | | | | | | | | | | | | | | | | | | | |
451 WVRFLLARHSGCRACVLGKDFRFLGKCPKKLSEGDGAELRKLQPSLNSG 500

501 ...HLASETLGDVGVPQHRGRGAWPRGSSLSSESSEGDFTLANTMEEILEE 547
| | | | | | | | : . | | | | | | | | | | | | | | | | | |
501 RLLHLAMRGLGELGAQPQQDHARWPRGSSLSECSEGDVTMANTMEEILEE 550

548 SEI 550
| | |
551 SEI 553

```

FIGURE 9



12/12

GL2R_RAT_R	MRPQSPSPVPSRCPREAPVPRVRAQPVGIPEAQGPVPLHSQQMRLLWGPG-RPFLALLLLV
HWBRPAT_TR	MKLGSSRAGPGRGSAGLLPGVHELPMGIPAPWGTSPLSFHRKCSLWAPG-RPFLTLVLLV
GLPR_HUMAN	-----MAGAPGPLRLALLLLGMVGRAGPRP-----
	* * *
GL2R_RAT_R	SIKQVTGSLLEKETTQKWANYKEKCLEDLHNRL---SGIFCNGTFFDRYVCWPHSPG-NVS
HWBRPAT_TR	SIKQVTGSLLEETTRKWAQYKQACLRDLLKEP---SGIFCNGTFFDQYVCWPHSSPG-NVS
GLPR_HUMAN	---QGATVSLWETVQKWREYRRCQORSLTEDPPATDLFCNRTFDEYACWPDGEPGSFVN
	* * * * *
GL2R_RAT_R	VPCPSYLPWNAESPGRAYRHCLAQGTWQTRENTTDIWQDESECSSENHSFRONVDHYALL
HWBRPAT_TR	VPCPSYLPWWESESSGRAYRHCLAQGTWQTIENTATDIWQDDSECSSENHSFKQNVDRYALL
GLPR_HUMAN	VSCPWYLPWASSVPQGHVYRFCTAEGWLQKDNSSLPRDLSECEESKRGRSSPEEQLL
	* * * * *
GL2R_RAT_R	YTLQLMYTVGYSVSLISLFLALTFLFLRKLHCTRNHYIHMNLFASFILKVLAVLVKDMVS
HWBRPAT_TR	STLQLMYTVGYSFSLISLFLALTLLFLRKLHCTRNHYIHMNLFASFILRTLAVLVKDVVF
GLPR_HUMAN	F-LYIIYTVGYALSFSALVIASAILLGFRHLHCTRNHYIHLNLFASFILRLSVFIKDAAL
	* * * * *
GL2R_RAT_R	HNSYSKRPDDESGWMSYLS-ETSVSCRSVQVLLHYFVGTNHLWLLVEGLYLHTLLEPTVF
HWBRPAT_TR	YNSYSKRPDNENGWMSYLS-EMSTSCRSVQVLLHYFVGANYLWLLVEGLYLHTLLEPTVL
GLPR_HUMAN	KWMYST-AAQHQWDGLLSYQDSLSCRLVFLMQYCVAAANYWLLVEGVLYTLTLLAFSVF
	* * * * *
GL2R_RAT_R	PERRLWPXYLVVGWAFPMFLFVVPWGFARAHLENTRCWATNGNLKIWWIIRGPMMLCVTVN
HWBRPAT_TR	PERRLWPXYLLLGWAFPVLFVVPWGFARAHLENTGCWTTNGNKKIWWIIRGPMMLCVTVN
GLPR_HUMAN	SEQWIFRLYVSIWGVPLLFVVPWGVIVKLYEDEGCWTRNSNMNYWLIIRLPILFAIGVN
	* * * * *
GL2R_RAT_R	FFIFLKILKLLISKLAHQMCFRDYKYRLAKSTLLLIPLLGVHEVLFTFFPDQVQGFSG
HWBRPAT_TR	FFIFLKILKLLISKLAHQMCFRDYKYRLAKSTLVLIPLLGVHEILFSFITDDQVEGFAK
GLPR_HUMAN	FLIFVRVICIVVSKLANLMCKTDIKCRLAKSTLTLLIPLGTHEVIFAFVMDHARGTLR
	* * * * *
GL2R_RAT_R	RIRLFIQLTLSSVHGFLVALQYGFANGEVKAELRKSWGRFLLARHWGCRTCVLGKNFRFL
HWBRPAT_TR	LIRLFIQLTLSSFHGFVALQYGFANGEVKAELRKYVVRFLLARHSGCRACVLGKDFRFL
GLPR_HUMAN	FIKLFTLSFTSFQGLMVAILYCFVNNEVQLEFRKSWERWRLEHLHIQDSSMKP-----
	* * * * *
GL2R_RAT_R	GKCSKILSEGDGSETLQKLRFTCSS---HLASETLGDVGVPHRGRGAWPRGSSLSSESS
HWBRPAT_TR	GKCPKILSEGDGAELRKLQPSLNSGRLLHLAMRGLGELGAQPPQDHARWPRGSSLSSECS
GLPR_HUMAN	LKCPTSS-LSSGATAGSSMYTATCQASCS-----
	* * *
GL2R_RAT_R	EGDFTLANTMEEILEESEI
HWBRPAT_TR	EGDVTMANTMEEILEESEI
GLPR_HUMAN	-----

FIGURE 10

CORRECTED  
VERSION\*CORRECT  
VERSION\*

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : C12N 15/12, C07K 14/705, C12N 5/10, 15/62, C07K 16/28, C12Q 1/68, G01N 33/50		A3	(11) International Publication Number: <b>WO 98/25955</b>
			(43) International Publication Date: 18 June 1998 (18.06.98)
(21) International Application Number: PCT/CA97/00969		(72) Inventors; and	
(22) International Filing Date: 15 December 1997 (15.12.97)		(75) Inventors/Applicants (for US only): MUNROE, Donald, G. [CA/CA]; 27 Wakefield Lane, Waterdown, Ontario L0R 2H3 (CA). GUPTA, Ashwani, K. [CA/CA]; 7031 Dunrobin Way, Mississauga, Ontario L5N 6Y4 (CA). VYAS, Tejal, B. [CA/CA]; 275 Riel Drive, Mississauga, Ontario L5B 3K1 (CA). McCALLUM, Kirk [CA/CA]; 3918 Pitch Pine Crescent, Mississauga, Ontario L5L 1P8 (CA). FAN, Ermei [CA/CA]; Apartment #418, 20 Southport Street, Toronto, Ontario M6S 4Y8 (CA).	
(30) Priority Data:			
08/767,224	13 December 1996 (13.12.96)	US	
08/787,721	24 January 1997 (24.01.97)	US	
08/845,546	24 April 1997 (24.04.97)	US	
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications			
US	08/767,224 (CIP)		
Filed on	13 December 1996 (13.12.96)		
US	08/787,721 (CIP)		
Filed on	24 January 1997 (24.01.97)		
US	08/845,546 (CIP)		
Filed on	24 April 1997 (24.04.97)		
(71) Applicant (for all designated States except US): ALLELIX BIOPHARMACEUTICALS INC. [CA/CA]; 6850 Goreway Drive, Mississauga, Ontario L4V 1V7 (CA).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
		<b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
		(88) Date of publication of the international search report: 30 July 1998 (30.07.98)	
(54) Title: CLONED GLUCAGON-LIKE PEPTIDE-2 RECEPTORS			
(57) Abstract			
<p>The invention relates to nucleotides and amino acid sequences encoding glucagon-like peptide 2 receptors, recombinant host cells transformed with such nucleotides, and methods of using the same in drug screening and related applications.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## CLONED GLUCAGON-LIKE PEPTIDE-2 RECEPTORS

### FIELD OF THE INVENTION

The invention is in the field of molecular biology. It relates, more particularly, to cloned glucagon-like peptide 2 receptors and their use in drug screening and related applications.

### BACKGROUND TO THE INVENTION

Glucagon-like peptide-2 (GLP-2) is a 33 amino acid peptide, which is expressed in a tissue determined fashion from the pleiotrophic glucagon gene and is highly related in terms of amino acid sequence to glucagon and Glucagon-like peptide-1 (GLP-1).

Mammalian forms of GLP-2 are highly conserved: for example, the human and degu (a south American rodent) forms differ by one and three amino acids respectively from rat GLP-2. Recently it was demonstrated that GLP-2 is an intestinotrophic peptide hormone; when given exogenously, GLP-2 can produce a marked increase in the proliferation of small intestinal epithelium of the test mice (Drucker et al, (1996) PNAS, 93:7911-7961). More recently, GLP-2 has been shown to increase D-Glucose maximal transport rate across the intestinal basolateral membrane (Cheeseman and Tseng: American Journal of Physiology (1996) 271:G477-G482).

To accelerate research into gastrointestinal biology and development of drugs useful in the treatment of various medical conditions including gastrointestinal disorders, it would be useful to provide the receptor through which the effects of GLP-2 are mediated.

### SUMMARY OF THE INVENTION

The GLP-2 receptor has now been cloned and characterized. Accordingly, the present invention provides an isolated polynucleotide encoding a GLP-2 receptor, particularly including mammalian forms and homologs thereof such as, in specific embodiments, the rat and human forms. In aspects of the invention, polynucleotide coding for a GLP-2 receptor is utilized for expression to obtain functional receptor protein and, in optionally labelled form, for further gene cloning to identify structurally related receptor proteins. In related aspects of the invention, anti-sense versions of GLP-2 receptor-encoding polynucleotides and fragments thereof are obtained and utilized to regulate GLP-2 receptor expression.

5 In another of its aspects, the invention provides GLP-2 receptor as a product of recombinant production in a cellular host. In related aspects, there are provided recombinant host cells that express GLP-2 receptor, as well as receptor-bearing membranes derived from such cells, and expression constructs in which polynucleotide coding for the GLP-2 receptor is linked to expression controls functional in the selected host cell.

10 In another of its aspects, the GLP-2 receptor is utilized in a chemicals screening program to identify GLP-2 receptor ligands. This method comprises the steps of incubating the candidate ligand with a GLP-2 receptor-producing cell of the present invention, or with a membrane preparation derived therefrom, and then measuring whether, or the extent to which, binding has occurred. Using cells that express a GLP-2 receptor coupled functionally to a second messenger system, such binding can be determined indirectly, to reveal ligand against activity, by detecting an appropriate reporter.

15 In another of its aspects, the invention provides antibodies directed to the GLP-2 receptor, for use for example in diagnostic procedures.

The invention is further described with reference to the following drawings in which:

20 BRIEF REFERENCE TO THE DRAWINGS

Figure 1 discloses a cDNA sequence (SEQ ID NO: 1), nucleotides 137-1789 of which encode the rat GLP-2 receptor wherein ambiguous base pairs are indicated using the standard IUB nomenclature (R: A or G, Y: C or T, M: A or C, K: G or T, S: G or C, W: A or T).

Figure 2 discloses the amino acid sequence of the expression product (SEQ ID NO: 2) from the cDNA of Figure 1.

30 Figure 3 illustrates the relative potencies of GLP-2 peptide and GLP-1 peptides for the receptor encoded by SEQ ID NO: 1.

Figure 4 discloses a cDNA sequence of 667 nucleotides (SEQ ID NO: 9) which encodes a 222 amino acid fragment (SEQ ID NO: 10) of a human GLP-2 receptor.

Figure 5 discloses the amino acid sequence (SEQ ID NO: 10) expressed from the cDNA of Figure 4.

Figure 6 discloses a cDNA sequence (SEQ ID NO: 11), nucleotides 121-1779 of which encode a human GLP-2 receptor.

Figure 7 discloses the amino acid sequence of the expression product (SEQ ID NO: 12) from the cDNA of Figure 6.

Figure 8 illustrates the functional activation by GLP-2 peptide of the human receptor encoded by SEQ ID NO: 11 (Figure 8).

Figure 9 compares the amino acid sequences of the rat GLP-2 receptor and the human GLP-2 receptor.

Figure 10 compares the amino acid sequences of the rat GLP-2 receptor and the human GLP-2 receptor against rat GLP-1 receptor.

#### DETAILED DESCRIPTION AND PREFERRED EMBODIMENTS

The invention relates in one respect to polynucleotides, in their isolated form, that code for GLP-2 receptors. As used herein "isolated" means separated from polynucleotides that encode other proteins. In the context of polynucleotide libraries, for instance, GLP-2 receptor-encoding polynucleotide is considered "isolated" when it has been selected, and hence removed from association with other polynucleotides within the library. Such polynucleotides may be in the form of RNA, or in the form of DNA including cDNA, genomic DNA and synthetic DNA. The GLP-2 receptors are characterized by structural features common to the G-protein coupled receptor class, including seven transmembrane regions, and by the functional properties of binding GLP-2 peptide selectively, i.e.,

preferentially, to GLP-1 peptide. Such selective binding is revealed as a statistically significant greater binding affinity of GLP-2 than of GLP-1, in the context of the assay chosen to measure such affinity. When expressed functionally in a host cell, i.e., in operable linkage with a responsive second messenger system the GLP-2 receptors are capable further of responding to GLP-2 binding by signal transduction. In this regard, the activity of a G-protein coupled receptor such as a GLP-2 receptor can be measured using any of a variety of appropriate functional assays in which activation of the receptor results in detectable change in the level of some second messenger system, such as adenylyl cyclase, calcium mobilization, inositol phospholipid hydrolysis products or guanylyl cyclase.

With reference to Figure 9 and Figure 10, which reveals homologies across amino acid sequences representing human and rat GLP-2 receptors, regions of 100% identity are indicated by solid vertical bars. In embodiments of the invention, the GLP-2 receptors are defined structurally as receptors that incorporate these regions of amino acid sequence and that also exhibit the functional characteristic of binding GLP-2 peptide selectively, relative to GLP-1 peptide. In more specific embodiments, the GLP-2 receptor structure further incorporates those amino acids which, across the human and rat receptor species, are highly conserved (indicated by ':'). At these sites, it will be appreciated that the sequence can contain any amino acid within the highly conserved amino acid family to which the identified amino acid belongs. In still more specific embodiments, the GLP-2 receptors have a structure which still further incorporates to the moderately conserved amino acids (indicated by '.') meaning, at these sites, that the amino acids within the moderately conserved family to which they belong. Beyond these sequences, the GLP-2 receptor structure can vary widely in embodiments of the invention, in allowing for non-conservative amino acid substitutions.

In one embodiment of the invention, the GLP-2 receptor is a rat GLP-2 receptor having the amino acid sequence of SEQ ID NO:2. In a specific embodiment of the invention, this rat GLP-2 receptor is encoded by the polynucleotide sequence of SEQ ID NO: 1. This particular GLP-2 receptor-encoding polynucleotide, also referred to as the WBR

gene, is a cDNA of rat origin. The expression product of this polynucleotide incorporates the mature form of the GLP-2 receptor, and additionally incorporates a secretion signal that is removed before membrane integration of the mature GLP-2 receptor product. Such a signal sequence may be naturally present on the polypeptides or replaced with a functionally equivalent secretion signal which is heterologous to the GLP-2 receptor. The replacement secretion signal chosen will depend on the expression system being used and will typically, but not essentially be endogenous to the chosen host and, also typically but not essentially, be homologous to the chosen expression controlling sequences.

The expressed rat GLP-2 receptor product (Figure 2, SEQ ID NO: 2) is characterized structurally as a single 550 amino acid polypeptide chain having a predicted molecular weight of 72kDa. Two functional translation start sites of the rat GLP-2 receptor have been identified, these are the codons encoding methionine 1 and methionine 42 of SEQ ID NO:2. Without wishing to be limited, by analogy with the GLP-1 receptor, it is believed that residues 1-66 of SEQ ID NO:2 are cleaved to provide a mature protein (i.e., the amino acid sequence of the receptor as it appears in the cell membrane) of 484 amino acids. With respect to structural domains of this GLP-2 receptor, hydropathy analysis and sequence alignment with related members of this sub-family of G protein coupled receptors indicates seven putative transmembrane domains, one spanning residues 181-203 inclusive (TM I), another spanning residues 211-230 (TM II), a third spanning residues 262-285 (TM III), a fourth spanning residues 300-321 (TM VI), a fifth spanning 339-362 (TM V), a sixth spanning 386-405 (TM VI) and a seventh spanning 422-441 (TM VII). Based on this assignment, it is likely that this GLP-2 receptor, in its natural membrane-bound form, consists of a an N-terminal extracellular domain, followed by a hydrophobic region containing seven transmembrane domains and an intracellular 442-550 amino acid C-terminal domain. The protein exhibits the highest degree of homology to the rat GLP-1 receptor with 49% identity at the amino acid level.

In a related embodiment, the GLP-2 receptor is of human origin (SEQ ID NO: 9) and incorporates the human GLP-2 receptor fragment having the amino acids of SEQ ID NO: 10.



This polynucleotide was isolated using the rat cDNA sequence as generally described below and as detailed in Example 3. In a further related embodiment of the invention, the cDNA is of human origin (SEQ ID NO: 11) and encodes the full length human GLP-2 receptor having residues 67-533 of the amino acid sequence of SEQ ID NO: 12. The human GLP-2 receptor precursor product (Figure 7, SEQ ID NO: 12) is characterized structurally as a single 553 amino acid polypeptide chain having a predicted molecular weight of 72 kDa. It is believed that, as for the rat GLP-2 receptor, this sequence, this precursor form of the human GLP-2 receptor incorporates an N-terminal signal sequence, which can be replaced by a functionally equivalent heterologous signal sequence. Without wishing to be limited it is believed that the mature form of the human GLP-2 receptor results after cleavage of residues 1-66 of SEQ ID NO:12 (Figure 7). With respect to structural domains of this GLP-2 receptor, hydropathy analysis and sequence alignment with related members of this sub-family of G protein coupled receptors indicates seven putative transmembrane domains, one spanning residues 181-203 inclusive (TM I), another spanning residues 211-230 (TM II), a third spanning residues 262-285 (TM III), a fourth spanning residues 300-321 (TM IV), a fifth spanning residues 339-362 (TM V), a sixth spanning residues 386-405 (TM VI), and a seventh spanning residues 422-441 (TM VII). Based on this assignment, it is likely that this GLP-2 receptor, in its natural membrane-bound form, consists of an N-terminal extracellular domain, followed by a hydrophobic region containing seven transmembrane domains interspersed with six short hydrophilic domains, and an intracellular domain, which is predicted to span residues 442-553. A second form of this GLP-2 receptor encompassed by the invention has a translational start site at the methionine codon at position 26 of the amino acid sequence presented in Figure 7 SEQ ID No. 12. The resulting 528 amino acid polypeptide chain also consists of an extracellular domain, seven transmembrane domains, and a C-terminal intracellular domain, and is at least 95% identical in sequence to residues 26-553 of the sequence presented in Figure 7, SEQ ID NO: 12.

In another embodiment, the invention provides GLP-2 receptor polynucleotide sequences and their unique sequence fragments as a tool useful to identify and isolate structurally related polynucleotides. At low stringency hybridization conditions, for instance, polynucleotide libraries can be probed to identify genes that are at least about 50% homologous to the GLP-

2 receptor gene. To facilitate isolation of rat GLP-2 receptor gene homologs that are also GLP-2 receptor-encoding, stringency conditions are desirably enhanced to identify homologs having at least 80% (medium stringency) sequence identity homology at the polynucleotide level to receptor gene. More desirably the WBR gene homologs are 90% identical, (high stringency) and most desirably they have at least 95% sequence (high stringency) identity when compared to WBR. Preferably, the isolated WBR homologs are characterized in that (1) they can be amplified using the PCR primers of SEQ ID NO: 3 and SEQ ID NO: 4 and (2) they bind to the probe of SEQ ID NO: 5 under high stringency conditions.

Still more preferably, the isolated homologs are those which bind, under conditions of high stringency, with consensus regions of the GLP-2 receptor-encoding polynucleotides, i.e., those regions of the rat and human GLP-2 receptor-encoding polynucleotides which are identical and, relative to GLP-1 receptor-encoding sequences, are also unique to the GLP-2 receptor-encoding polynucleotides. Alignment of this nature reveals a number consensus regions for GLP-2: for example nucleotides spanning from 1460-1786. In one embodiment of the invention, these sequences, and their complements, constitute polynucleotide fragments useful, as just described for the intact gene, to identify polynucleotides structurally related to the human and rat GLP-2 receptor embodiments of the invention.

In a related embodiment, the cDNA sequence or the unique fragments of the GLP-2 receptor, most preferably of the human GLP-2 receptor, can be used in appropriately labelled form, e.g.,  $p^{32}$  labelled, for diagnosis of conditions associated with aberrant expression of the GLP-2 receptor. For example, over or under expression or expression in an inappropriate tissue. In one embodiment appropriate PCR primers (e.g., regions unique to the GLP-2 receptor but conserved between species) can be used diagnostically to identify aberrant structure or levels of GLP-2 receptor mRNA, e.g., associated with inherited or acquired disease states.

It has been found that the human GLP-2 receptor is located on chromosome 17P13. Thus, in a further embodiment, the invention provides expression products from this locus that

hybridize with the human GLP-2 receptor polynucleotide(Figure 6; SEQ ID NO:11) under stringent conditions.

As source material to isolate GLP-2 receptor-encoding homologs of the rat GLP-2 receptor-encoding polynucleotide gene, it is desirable but not necessary to use libraries of fetal or mature hypothalamal, jejunal, hindbrain or stomach tissue obtained from the vertebrate species targeted for receptor isolation. The invention accordingly includes not only the rat GLP-2 receptor-encoding polynucleotide of SEQ ID NO:1, but structural homologs thereof and particularly those that code for proteins having GLP-2 receptor properties. As exemplified hereinbelow, the WBR gene has been used successfully as a starting material to clone the human homolog of the rat GLP-2 receptor. Thus, the invention provides polynucleotides that encode GLP-2 receptors, including rat GLP-2 receptor and vertebrate homologs, particularly mammalian homologs thereof including human homologs, as well as synthetic variants of these.

It will be appreciated that such homologs can also be identified in libraries by screening, as noted, with fragments of the rat receptor gene or the human homolog, which incorporate at least 15 nucleotides, and preferably at least 25 nucleotides. With reference to SEQ ID NO: 1 and the nucleotide numbering appearing thereon, suitable nucleotide fragments include in addition to the consensus nucleotide fragments noted above, those corresponding in sequence to the extracellular GLP-2 binding domain, and the stipulated transmembrane regions and the C-terminal portion of the receptor.

Technically, the identification of GLP-2 receptor genes can be achieved by applying standard hybridization or amplification techniques to a tissue-derived polynucleotide library. A wide variety of such libraries are commercially available. Where construction of a cDNA library is necessary, established techniques are applied. For example, isolation of such a WBR homolog typically will entail extraction of total messenger RNA from a fresh source of tissue, such as hypothalamal, jejunal, stomach or hindbrain tissue, preferably hypothalamal tissue or cell lines derived from these tissues, followed by conversion of message to cDNA and formation of a library in for example a bacterial plasmid, more typically a bacteriophage.

Such bacteriophage harboring fragments of the DNA are typically grown by plating on a lawn of susceptible E. coli bacteria, such that individual phage plaques or colonies can be isolated. The DNA carried by the phage colony is then typically immobilized on a nitro-cellulose or nylon-based hybridization membrane, and then hybridized, under carefully controlled conditions, to a radioactively (or otherwise) labelled probe sequence to identify the particular phage colony carrying the fragment of DNA of particular interest, in this case a rat or human GLP-2 homolog. The phage carrying the particular gene of interest is then purified away from all other phages from the library, in order that the foreign gene may be more easily characterized. Typically, the gene or a portion thereof is then isolated by subcloning into a plasmidic vector for convenience, especially with respect to the full determination of its DNA sequence.

As an alternative to obtaining GLP-2 encoding DNA directly as a DNA insert from an available or a constructed cDNA library, in light of the present disclosure it can be synthesized de novo using established techniques of gene synthesis. Because of the length of the GLP-2 receptor-encoding DNAs of SEQ ID NO: 1, SEQ ID NO: 9 and SEQ ID NO: 11, application of automated synthesis may require staged gene construction, in which regions of the gene up to about 300 nucleotides in length are synthesized individually and then ligated in correct succession for final assembly. Individually synthesized gene regions can be amplified by PCR. The application of automated synthesis may typically be applied by synthesizing specific regions or fragments of the gene and ligating them, usually via designed overlaps, in correct succession to form the final gene sequence. In this case, the longer the oligonucleotide building blocks, the fewer will be the ligations needed, resulting in greater ease of assembly.

The application of automated gene synthesis techniques provides an opportunity for generating sequence variants of the naturally occurring GLP-2 receptor gene. It will be appreciated, for example, that polynucleotides coding for the GLP-2 receptor herein described can be generated by substituting one or more synonymous codons for those represented in the naturally occurring polynucleotide sequences herein provided and such "synonymous codon equivalents" are within the scope of the present invention. In addition, polynucleotides

coding for synthetic variants of the GLP-2 receptor herein provided can be generated which incorporate from 1 to 20, e.g., from 1 to 5, amino acid substitutions, or deletions or additions. Preferred sites for such a modification include areas of non homology between the rat and human sequence, for example amino acid ranges 70-92, 328-350 and 475-504. Since it will be desirable typically to retain the natural ligand binding profile of the receptor for screening purposes, it is desirable to limit amino acid substitutions, for example to the so-called conservative replacements in which amino acids of similar charge are substituted (Fig. 9), and to limit substitutions to those sites less critical for receptor activity. For example, substitution of nucleotides "G" and "A" for nucleotides "A" and "G" respectively at positions 374 and 375 of the human cDNA sequence of SEQ ID NO: 11; resulting in the replacement of the naturally occurring arginine residue at position 85 of SEQ ID NO: 12 with a glutamic acid residue, provides a functional receptor. This functional receptor is referred to herein as the Glu<sup>85</sup> variant human GLP-2 receptor.

Having obtained GLP-2 receptor encoding polynucleotide, GLP-2 receptor can be produced in a number of ways, including in vitro transcription and via incorporation of the DNA into a suitable expression vector and expression in the appropriate host, for example in a bacterium such as E.coli, in yeast or in insect or in a mammalian cell. A variety of gene expression systems have been adapted for use with these hosts and are now commercially available, and any one of these systems can be selected to drive expression of the GLP-2 receptor-encoding DNA. Expression vectors may be selected to provide transformed cell lines that express the receptor-encoding DNA either transiently or in a stable manner. For transient expression, host cells are typically transformed with an expression vector harboring an origin of replication functional in a mammalian cell. For stable expression, such replication origins are unnecessary, but the vectors will typically harbor a gene coding for a product that confers on the transformants a survival advantage, to enable their selection such as a gene coding for neomycin resistance in which case the transformants are plated in medium supplemented with neomycin.

These expression systems, available typically but not exclusively in the form of plasmidic vectors, incorporate expression cassettes the functional components of which include DNA

constituting expression controlling sequences and optionally also signal peptides encoding sequences, which are host-recognized and enable expression of the receptor-encoding DNA when linked 5' thereof. The systems further incorporate DNA sequences which terminate expression when linked 3' of the receptor-encoding region. Thus, for expression in the selected mammalian cell host, there is generated a recombinant DNA expression construct in which the receptor-encoding DNA is linked with expression controlling DNA sequences recognized by the host, and which include a region 5' of the receptor-encoding DNA to drive expression, and a 3' region to terminate expression.

Included among the various recombinant DNA expression systems that can be used to achieve mammalian cell expression of the receptor-encoding DNA are those that exploit promoters of viruses that infect mammalian cells, such as the promoter from the cytomegalovirus (CMV), the Rous sarcoma virus (RSV), simian virus (SV40), murine mammary tumor virus (MMTV) and others. Also useful to drive expression are promoters such as the LTR of retroviruses, insect cell promoters such as those regulated by temperature, and isolated from *Drosophila*, as well as mammalian gene promoters such as those regulated by heavy metals i.e. the metallothionein gene promoter, and other steroid-inducible promoters.

In another of its aspects, the invention provides cells or membranes derived therefrom which are adapted by genetic alteration for use, for example, in identifying GLP-2 receptor ligands. In preferred embodiments, such cells are adapted genetically by the insertion of polynucleotide coding for a GLP-2 receptor. In particularly preferred embodiments, such cells incorporate a recombinant DNA molecule, e.g. an expression construct/vector, in which DNA coding for the GLP-2 receptor and expression controlling elements functional in the host are linked operably to drive expression of the DNA. For incorporation of receptor into cell plasma membranes, the vector can, if desired, be designed to provide a suitable heterologous signal peptide sequence to substitute for the signal peptide encoded naturally within the receptor DNA.

Suitable GLP-2 producing cells include the Chinese hamster ovary (CHO) cells for example of K1 lineage (ATCC CCL 61) including the Pro5 variant (ATCC CRL 1281); the fibroblast-like cells derived from SV40-transformed African Green monkey kidney of the CV-1 lineage (ATCC CCL 70), of the COS-1 lineage (ATCC CRL 1650) and of the COS-7 lineage (ATCC CRL 1651); murine L-cells, murine 3T3 cells (ATCC CRL 1658), murine C127 cells, human embryonic kidney cells of the 293 lineage (ATCC CRL 1573), human carcinoma cells including those of the HeLa lineage (ATCC CCL 2), and neuroblastoma cells of the lines IMR-32 (ATCC CCL 127), SK-N-MC (ATCC HTB 10) and SK-N-SH (ATCC HTB 11).

For use in ligand screening assays, cell lines expressing the receptor-encoding DNA can be stored frozen for later use. Such assays may be performed either with intact cells, or with membrane preparations derived from such cells. The membrane preparations typically provide a more convenient substrate for the ligand binding experiments, and are therefore preferred as binding substrates. To prepare membrane preparations for screening purpose, i.e., ligand binding experiments, frozen intact cells are homogenized while in cold water suspension and a membrane pellet is collected after centrifugation. The pellet is then washed in cold water, and dialyzed to remove any endogenous GLP-2 receptor ligands that would otherwise compete for binding in the assays. The dialyzed membranes may then be used as such, or after storage in lyophilized form, in the ligand binding assays.

The binding of a candidate ligand to a selected GLP-2 receptor of the invention can be assessed typically using a predetermined amount of cell-derived membrane (measured for example by protein determination), generally from about 25ug to 100ug. Generally, competitive binding assays will be useful to evaluate the affinity of a test compound relative to GLP-2. This competitive binding assay is performed by incubating the membrane preparation with radiolabelled GLP-2 peptide, for example [ $H^3$ ] or a radioiodinated GLP-2 analog, in the presence of unlabelled test compound added at varying concentrations. Following incubation, either displaced or bound radiolabelled GLP-2 can be recovered and measured, to determine the relative binding affinities of the test compound and GLP-2 for the

GLP-2 receptor used as substrate. In this way, the affinities of various compounds for the GLP-2 receptor can be measured.

Alternatively, binding of a candidate ligand to a GLP-2 receptor can be assessed using a functional assay. Using this approach, for example, intact cells harvested about two days following transient transfection or after about the same period following plating of stably transfected cells can be used to assess ligand binding. In a preferred embodiment, 293 EBNA cells (Invitrogen Cat. R620-07) are stably transformed with the pREP7 vector (Invitrogen Cat. V007-50) incorporating expressibly therein a GLP-2 receptor. Thereafter, binding of an agonist (or using a competition base format an antagonist) to the receptor can be discerned by measuring the level of intracellular cAMP. Most conveniently, intracellular cAMP is measured indirectly using a reporter system, wherein an easily measurable and preferably easily quantifiable downstream event indicates the level of intracellular cAMP. For example, measuring the level of the expression of a reporter gene construct having polynucleotide sequence under the control of a promoter which is responsive to cAMP. Alternatively, measurement of intracellular calcium, released from intracellular stores in response to an increase in intracellular cAMP, can be used as an indicator of the level of intracellular cAMP, for example, by incorporating into the transformed cell a protein that fluoresces on binding to calcium. In a preferred embodiment, intracellular cAMP levels are measured using the commercially available EIA kit. An additional advantage of the functionally based approach to assessing ligand binding is that the system can be automated allowing high throughput and ultra high through screening of vast chemical libraries.

As an alternative to using cells that express receptor-encoding DNA, ligand characterization may also be performed using cells for example *Xenopus* oocytes, that yield functional membrane-bound receptor following introduction of messenger RNA coding for the GLP-2 receptor. In this case, the GLP-2 receptor-encoding polynucleotide of the invention is typically subcloned into a plasmidic vector such that the introduced gene may be easily transcribed into RNA via an adjacent RNA transcription promoter supplied by the plasmidic vector, for example the T3 or T7 bacteriophage promoters. RNA is then transcribed from the inserted gene in vitro, and can then be injected into *Xenopus* oocytes. Each oocyte is a



single cell, but is large enough to be penetrated by a fine-tipped microneedle without causing irreparable damage. Following the injection of nL volumes of an RNA solution, the oocytes are left to incubate for up to several days, whereupon the oocytes are tested for the ability to respond to a particular ligand molecule supplied in a bathing solution.

5  
Candidate GLP-2 receptor ligands can vary widely in structure, and most suitably include proteins which are highly related to GLP-2 itself in terms of amino acid sequence. For instance, the peptides disclosed in co-pending United States patent applications WO97/39031 and WO96/32414, incorporated herein by reference, may usefully be screened for GLP-2  
10 receptor binding activity.

In addition to naturally occurring GLP-2 receptor sequences functional chimeric, receptors, incorporating portions of the GLP-2 receptor sequence and the polynucleotides encoding them are also embodiments of the invention. Functional chimeric GLP-2 receptors are constructed  
15 by combining the extracellular receptive sequences of a GLP-2 receptor with one or more of the transmembrane and intracellular segments of a known seven transmembrane G-protein coupled receptors for test purposes. This concept was demonstrated by Kobilka et al. (1988, Science 240:1310-1316) who created a series of chimeric  $\alpha 2$ - $\beta 2$  adrenergic receptors (AR) by inserting progressively greater amounts of  $\alpha 2$ -AR transmembrane sequence into  $\beta 2$ -AR. The  
20 binding activity of known agonists changed as the molecule shifted from having more  $\alpha 2$  than  $\beta 2$  conformation, and intermediate constructs demonstrated mixed specificity. The specificity for binding antagonists, however, correlated with the source of the transmembrane domain VII. The importance of transmembrane domain VII for ligand recognition was also found in chimeras utilizing two yeast  $\alpha$ -factor receptors and is significant because the yeast  
25 receptors are classified as miscellaneous receptors. Thus, functional role of specific domains appears to be preserved throughout the seven transmembrane G-protein coupled receptor family regardless of category.

In parallel fashion, internal segments or cytoplasmic domains from a particular GLP-2  
30 receptor are exchanged with the analogous domains of a known seven transmembrane G-protein coupled receptor and used to identify the structural determinants responsible for coupling the

receptors to trimeric G-proteins (Dohlman et al. (1991) *Annu Rev Biochem* 60:653-688). A chimeric receptor in which domains V, VI, and the intracellular connecting loop from  $\beta 2$ -AR were substituted into  $\alpha 2$ -AR was shown to bind ligands with  $\alpha 2$ -AR specificity, but to stimulate adenylate cyclase in the manner of  $\beta 2$ -AR. This demonstrates that for adrenergic-type receptors, G-protein recognition is present in domains V and VI and their connecting loop. The opposite situation was predicted and observed for a chimera in which the V->VI loop from  $\alpha 1$ -AR replaced the corresponding domain on  $\beta 2$ -AR and the resulting receptor bound ligands with  $\beta 2$ -AR specificity and activated G-protein-mediated phosphatidylinositol turnover in the  $\alpha 1$ -AR manner. Finally, chimeras constructed from muscarinic receptors also demonstrated that V->VI loop is the major determinant for specificity of G-protein activity.

Chimeric or modified seven transmembrane G-protein coupled receptors containing substitutions in the extracellular and transmembrane regions have shown that these portions of the receptor determine ligand binding specificity. For example, two Ser residues conserved in domain V of all adrenergic and D catecholamine receptors are necessary for potent agonist activity. These serines are believed to form hydrogen bonds with the catechol moiety of the agonists within the binding site. Similarly, an Asp residue present in domain III of all seven transmembrane G-protein coupled receptors which bind biogenic amines is believed to form an ion pair with the ligand amine group in the binding site.

Functional, cloned seven transmembrane G-protein coupled receptors are expressed in heterologous expression systems and their biological activity assessed (e.g., Marullo et al. (1988) *Proc Natl Acad Sci* 85:7551-7555; King et al. (1990) *Science* 250:121-123). One heterologous system introduces genes for a mammalian seven transmembrane G-protein coupled receptors and a mammalian G-protein into yeast cells. The seven transmembrane G-protein coupled receptor is shown to have appropriate ligand specificity and affinity and trigger appropriate biological activation--growth arrest and morphological changes--of the yeast cells.

An alternate procedure for testing chimeric receptors is based on the procedure utilizing the  $P_{2u}$  purinergic receptor ( $P_{2u}$ ) as published by Erb et al. (1993, *Proc Natl Acad Sci* 90:104411-

104453). Function is easily tested in cultured K562 human leukaemia cells because these cells lack  $P_{2u}$  receptors. K562 cells are transfected with expression vectors containing either normal or chimeric  $P_{2u}$  and loaded with fura-a, fluorescent probe for  $Ca^{++}$ . Activation of properly assembled and functional  $P_{2u}$  receptors with extracellular UTP or ATP mobilizes intracellular  $Ca^{++}$  which reacts with fura-a and is measured spectrofluorimetrically. As with the seven transmembrane G-protein coupled receptors above, chimeric genes are created by combining sequences for extracellular receptive segments of any newly discovered seven transmembrane G-protein coupled receptors polypeptide with the nucleotides for the transmembrane and intracellular segments of the known  $P_{2u}$  molecule. Bathing the transfected K562 cells in microwells containing appropriate ligands triggers binding and fluorescent activity defining effectors of the seven transmembrane G-protein coupled receptors molecule. Once ligand and function are established, the  $P_{2u}$  system is useful for defining antagonists or inhibitors which block binding and prevent such fluorescent reactions.

In addition to using the receptor-encoding DNA to construct cell lines useful for ligand screening, expression of the DNA can according to another aspect of the invention be performed to produce fragments of the receptor in soluble form, for structure investigation, to raise antibodies and for other experimental uses. It is expected that the extracellular portion of the GLP-2 receptor contributes significantly to the binding of ligand molecule. It is therefore desirable in the first instance to facilitate the characterization of the receptor-ligand interaction by providing this extracellular ligand-binding domain in quantity and in isolated form, i.e., free from the remainder of the receptor. Such a construct has been made for the rat GLP-1 receptor, and it was shown to bind GLP-1 (Wilmen et al. (1996) FEBS LETTS, 398:43-47).

To accomplish this, the full-length GLP-2 receptor-encoding DNA may be modified by site-directed mutagenesis, so as to introduce a translational stop codon into the extracellular N-terminal region, immediately before the sequence encoding the first transmembrane domain (TM1), i.e., before residue 181 of SEQ ID NO: 2 and before residue 181 of SEQ ID NO: 12. Since there will no longer be produced any transmembrane domain(s) to "anchor" the receptor into the membrane, expression of the modified gene will result in the secretion, in

soluble form, of only the extracellular ligand-binding domain. Standard ligand-binding assays may then be performed to ascertain the degree of binding of a candidate compound to the extracellular domain so produced. It may of course be necessary, using site-directed mutagenesis, to produce several different versions of the extracellular regions, in order to optimize the degree of ligand binding to the isolated domains.

It will be appreciated that the production of such extracellular ligand binding domains may be accomplished in a variety of host cells. Mammalian cells such as CHO cells may be used for this purpose, the expression typically being driven by an expression promoter capable of high-level expression, for example the CMV (cytomegalovirus) promoter. Alternately, non-mammalian cells, such as insect Sf 9 (*Spodoptera frugiperda*) cells may be used, with the expression typically being driven by expression promoters of the baculovirus, for example the strong, late polyhedron protein promoter. Filamentous fungal expression systems may also be used to secrete large quantities of such extracellular domains of the GLP-2 receptor.

*Aspergillus nidulans*, for example, with the expression being driven by the *alcA* promoter, would constitute such an acceptable system. In addition to such expression hosts, it will be further appreciated that any prokaryotic or other eukaryotic expression system capable of expressing heterologous genes or gene fragments, whether intracellularly or extracellularly would be similarly acceptable.

The availability of isolated extracellular ligand-binding domains of the receptor protein makes it feasible to determine the 3-dimensional structures of these ligand-binding regions, with or without a candidate ligand complexed thereto, by a combination of X-ray crystallographic and advanced 2D-NMR techniques. In this way, additional new candidate compounds, predicted to have the required interactions with the 3-dimensional receptor structure, can be specifically designed and tested.

With large domains, crystallography is the method of choice for structure determination of both the domain in isolation, and of the co-complex with the natural ligand (or an appropriate antagonist or agonist molecule). If a particular domain can be made small enough, for example approximately 100-130 amino acids in length, then the powerful technique of 2-D

NMR can also be applied to structure determination. This enables not only the determination of the domain structure, but also provides dynamic information about the drug-receptor interaction.

5 For use particularly in detecting the presence and/ or location, for example in intestinal tissue, the present invention also provides, in another of its aspects, labelled antibody to a GLP-2 receptor. To raise such antibodies, there may be used as immunogen either the intact, soluble receptor or an immunogenic fragment thereof, produced in a microbial or mammalian cell host as described above or by standard peptide synthesis techniques. Regions of the GLP-2  
10 receptor particularly suitable for use as immunogenic fragments include those corresponding in sequence to an extracellular region of the receptor, or a portion of the extracellular region, such as peptides consisting of 10 or more amino acids of the 401-509 region of SEQ ID NO: 2. With regard to the human GLP-2 receptor (SEQ ID NO: 12), peptides comprising the mature extracellular domain (residues 65-180); intracellular loop 3 (residues 363-385) and the  
15 intracellular C-terminal domain (residues 442-533) may be usefully employed as immunogens for the production of antibodies to the human GLP-2 receptor.

Antibodies to the desired GLP-2 receptor or fragment immunogen are available, for polyclonal antibody production, from the blood of an animal that has been immunized with  
20 the immunogen. Alternatively, for monoclonal antibody production, immunocytes such as splenocytes can be recovered from the immunized animal and fused, using hybridoma technology, to a myeloma cells. The fusion products are then screened by culturing in a selection medium, and cells producing antibody are recovered for continuous growth, and antibody recovery. Recovered antibody can then be coupled covalently to a detectable label,  
25 such as a radiolabel, enzyme label, luminescent label or the like, using linker technology established for this purpose.

Animal model systems which elucidate the physiological and behavioral roles of the GLP-2 receptor are produced by creating transgenic animals in which the activity of the  
30 GLP-2 receptor is either increased or decreased, or the amino acid sequence of the expressed GLP-2 receptor is altered, by a variety of techniques. Examples of these

techniques include, but are not limited to: 1) Insertion of normal or mutant versions of DNA encoding a GLP-2 receptor, by microinjection, electroporation, retroviral transfection or other means well known to those skilled in the art, into appropriate fertilized embryos in order to produce a transgenic animal or 2) Homologous recombination of mutant or normal, human or animal versions of these genes with the native gene locus in transgenic animals to alter the regulation of expression or the structure of these GLP-2 receptor sequences. The technique of homologous recombination is well known in the art. It replaces the native gene with the inserted gene and so is useful for producing an animal that cannot express native GLP-2 receptors but does express, for example, an inserted mutant GLP-2 receptor, which has replaced the native GLP-2 receptor in the animal's genome by recombination, resulting in under expression of the transporter. Microinjection adds genes to the genome, but does not remove them, and so is useful for producing an animal which expresses its own and added GLP-2 receptors, resulting in over expression of the GLP-2 receptors.

One means available for producing a transgenic animal, with a mouse as an example, is as follows: Female mice are mated, and the resulting fertilized eggs are dissected out of their oviducts. The eggs are stored in an appropriate medium such as M2 medium. DNA or cDNA encoding a GLP-2 receptor is cesium chloride purified from a vector by methods well known in the art. Inducible promoters may be fused with the coding region of the DNA to provide an experimental means to regulate expression of the transgene. Alternatively or in addition, tissue specific regulatory elements may be fused with the coding region to permit tissue-specific expression of the trans-gene. The DNA, in an appropriately buffered solution, is put into a microinjection needle (which may be made from capillary tubing using a piper puller) and the egg to be injected is put in a depression slide. The needle is inserted into the pronucleus of the egg, and the DNA solution is injected. The injected egg is then transferred into the oviduct of a pseudopregnant mouse ( a mouse stimulated by the appropriate hormones to maintain pregnancy but which is not actually pregnant), where it proceeds to the uterus, implants, and develops to term. As noted above, microinjection is not the only methods for inserting DNA into the egg cell, and is used here only for exemplary purposes.

The invention having been described above, may be better understood by referring to the following examples. The following examples are offered for the purpose of illustrating the invention and should not be interpreted as a limitation of the invention.

## EXAMPLE 1

### Isolation of the GLP-2 receptor

#### PCR-assisted cloning of partial rat and mouse GLP-2 receptor cDNAs

Rat Neonate Intestine cDNA library (Stratagene, La Jolla, CA; Cat. 936508) and Mouse Jejunum first strand cDNA was prepared. Degenerate primers M-2F/S (SEQ ID NO: 3) and M-7R/S (SEQ ID NO: 4) were used to amplify a partial fragment of the rat GLP-2 receptor from the Rat Neonate Intestine cDNA library and of the mouse GLP-2 receptor from Mouse Jejunum template. The protocol is described below:

#### Degenerate PCR :

6 µl 10x VENT buffer from New England Biolabs

6 µl 2.5µM each stock dATP, dCTP, dGTP and dTTP

4 µl rat neonate intestine cDNA (1:10 dilution)

3 µl 25µM M2F/S primer [5'-TTTTTCTAGAASRTSATSTACACNGTSGGCTAC-3']  
(SEQ ID NO: 3)

3 µl 25µM M7R/S primer [5'-TTTTCTCGAGCCARCARCCASSWRTARTTGGC-3']  
(SEQ ID NO: 4)

2 µl (10 units) Amplitaq DNA polymerase (Perkin Elmer)

36 µl ddH<sub>2</sub>O.

Reaction conditions: 35 cycles at 94°C, 2°min.; 94°C, 1 min.; 53°C, 30 sec.; 72°C, 1 min.

The predominant PCR product was a 303 base pair (bp) DNA fragment. 30 µl samples of the above PCR were purified using the QIAGEN PCR purification kit and eluted in 30 µl ddH<sub>2</sub>O. The resulting product was then re-amplified using the same degenerate PCR conditions, with the exception only of 31 cycles at 94°C.

The predominant product at 303 base pair (bp) was cut out and purified using QIAGEN QIAquick gel purification protocol into 30 µl ddH<sub>2</sub>O. The resulting product was then re-amplified using the same degenerate PCR conditions, with the exception only of 31 cycles at 94°C.

Next, double digest (Xba I and Xho I) was done on the entire reamplified PCR reaction as follows: 28 µl DNA; 16 µl 10X One-Phor-All buffer (Pharmacia); 2 µl (40 units) Xba I enzyme (Pharmacia); 2 µl (40 units) Xho I enzyme (Pharmacia); and 30 µl ddH<sub>2</sub>O.

The samples were digested 4 hours in 37°C water block heater, brought up to 100 µl volume with ddH<sub>2</sub>O (sterile) and purified by (1) equal amount (100ul) chloroform extraction; (2) weekend precipitation with 2 volumes ethanol/10 volumes 3M sodium acetate; (3) 1x wash with 70% EtOH; and (4) resuspension in 10 µl 1x TE (pH 8.0).

pBluescript clone 5HT1F#9 was next digested with Xba I and Xho I as follows:

10 µl DNA (pBluescript clone 5HT1F#9)  
5 µl 10X NEBuffer 2 (New England Biolabs)  
3 µl (1:20 dilution=3 units) Xba I (New England Biolabs)  
3 µl (1:20 dilution=3 units) Xho I (New England Biolabs)  
5 µl (10x) BSA (New England Biolabs)  
24 µl ddH<sub>2</sub>O.

The sample was digested for 3 hours in 37 C water block heater, heat-inactivated at 65°C for 20 min and purified using GeneCleanII kit from BIO 101. Aliquots of the PCR reactions were cloned into the above pBluescript plasmid vector using T4 DNA ligase kit (New England Biolabs) and transformed into Epicurean Coli XL-2 Blue MRF' Ultracompetent cells (Stratagene). The transformation was plated onto 2xYT + AMP plates and single colonies were picked. DNA minipreps were made using QIAGEN QIA-prep 8 miniprep kit and the sequences of the fragments were determined using ABI system. Novel sequences were identified containing a partial fragment of the rat and mouse GLP-2 receptor sequence.



Cloning of cDNA with complete GLP-2 receptor coding region was achieved as follows:  
First, cDNA libraries from the following three tissues were used for screening,

1. Rat Hypothalamus (RHT)
2. Rat Hind Brain (RHB)
- 5 3. Rat Duodenum and Jejunum (RDJ)

The three cDNA libraries were prepared by priming with random primer and subcloning unidirectionally into Hind III and Not I sites of pcDNA3.

Next, the three cDNA libraries were homology screened by a degenerate oligo C4-4  
10 [5'-AACTACATCCACMKGMAYCTGTTYVYGTCTTCATSCT-3'] (SEQ ID NO: 5)  
by colony lifts and filter hybridization. The following hybridization conditions were  
employed: 5X SSPE (1X SSPE is 0.18M NaCl, 10mM NaH<sub>2</sub>PO<sub>4</sub> (pH 7.4), 10mM EDTA  
(pH 7.4)) and 5X Denharts solution (1% Ficoll, 1% Polyvinylpyrrolidone, 1% BSA); 25  
mg/ml salmon sperm DNA.

15 Filters were hybridized at 50°C overnight. Then the filters were washed 2 times in 2X  
SSPE and 1% SDS at room temperature for 30 min, 2 times in 2X SSPE and 1% SDS at  
50°C for 20 min per wash, and finally two times in 1X SSPE and 0.5% SDS. Positive  
clones were identified by autoradiography. A plug of 1cm<sup>2</sup> surrounding the positive  
20 clone was removed from the plate and placed in 1 ml of 2x YT +20% Glycerol, vortexed  
and was frozen at -80°C.

Plasmid DNA from positive plugs was prepared as follows: 100 ml of bacterial culture  
of each positive plug was grown on an agar plate. The bacterial cells were scraped and  
25 resuspended in 1 ml of 2xYT medium+20% Glycerol. Bacterial pellet from the 250 ml  
of bacterial resuspension was resuspended in 150 ml of Solution I (50mM Glucose,  
10mM Tris-HCl, 1mM EDTA), lyse in Solution II (0.2M NaOH, 1% SDS), neutralized  
with ice cold Solution III (Potassium acetate; 4 vol.of 5M potassium acetate + 1 vol. of  
10M acetic acid). After pelleting bacterial DNA, 340 ml isopropanol was added to the  
30 supernatant. This was centrifuged at max for 15 min. The pellet was resuspended in TE  
+ 20 mg/ml RNase, incubated at 37 C for 30 minutes and precipitated with isopropanol

+ 0.2M potassium acetate. After centrifugation, the pellet was washed with 70% alcohol, allowed to air dry and resuspended in TE.

Plasmid DNA from 2777-clone pools of rat hypothalamus cDNA library RHT cDNA library was next exploited as follows: Two primers were designed from an area of the PCR-cloned GLP-2 receptor cDNA sequence that did not have identity to known receptors of the gene family. The two primers P23-R1 and P23-F1 amplified a 196 bp fragment only from novel clone DNA but not with GLP-1 receptor cDNA or PACAP receptor cDNA. The Expand™ PCR system from Boehringer Mannheim (Cat. 1681-842) was used under the following conditions:

2 µl of 10x Expand™ Buffer 1

2.8 µl of 2.5mM dNTP mix

0.6 µl of primer P23-R1 [5'-TCATCTCCCTCTTCTTGGCTCTTAC-3'] (SEQ ID NO: 6)

0.6 µl of primer P23-F1 [5'-TCTGACAGATATGACATCCATCCAC-3'] (SEQ ID NO: 7)

0.3 µl of Expand PCR enzyme (1 unit)

12.7 µl water

1 µl DNA

Reaction conditions: 32 cycles at 93°C, 40 sec; (cycles) 58°C, 40 sec; (cycles) 68°C, 40 sec

DNAs from each positive plug or pool of 2777-clone pools were amplified with specific primers P23-F1 and P23-R1 under the conditions specified above. Out of 1057 C4-4 hybridization-positive plugs and 884 2777-clone pools only five template sources amplified a 196 bp PCR product. These were: (1) Plug 334, (2) Plug 780, (3) RHT pool 233, (4) RHT pool 440, and (5) RHT pool 587.

Amplification of GLP-2R cDNA from the five positive templates was then performed. By using one specific primer (P23-R1 or P23-F1) and one primer based on pcDNA3 vector (Invitrogen) sequence (830F or 1186R), the GLP-2R cDNA insert was directly

amplified from clonally impure plugs or 2777-clone pools. The sequences of the vector primers were as follows:

830F: [5'-AACCCACTGCTTAC-3']

1186R: [5'-CCCAGAATAGAATGACACC-3']

5 The PCR was done, under the following conditions just noted, using Expand™ PCR system.

10 The most prominent band was reamplified, purified and sequenced. Based on the amplified sequence obtained, additional primers were designed and new sequencing carried out. In this manner the complete sequences of the GLP-2R cDNA inserts in all five sources of clones were determined. Sequence analysis showed that only pool RHT 440 and pool RHT 587 contain clones with complete coding sequence of GLP-2R and that the two clones were identical (derived from the same cDNA clone).

15 Because of difficulty in clonally purifying the GLP-2 receptor cDNA clone from the RHT 440 or RHT 587 cDNA library pools, the cDNA was amplified and recloned into pcDNA3. Based on the sequence obtained from RHT 440 and RHT 587, two primers were designed one which primed starting 4bp upstream of the initiation codon and another which primed starting 8bp downstream of the stop codon.

20 WBR-C5: [5'-CAGGGGCCGGTACCTCTCCACTCC-3']

WBR-C3: [5'-TTGGGTCCTCGAGTGGCCAAGCTGCG-3']

25 The two primers were used to amplify a DNA fragment of approximately 1525 bp fragment under the following PCR conditions using Expand™ PCR system from Boehringer Mannheim (Catalogue no. 1681-842).

10 µl of 10x Expand™ PCR Buffer 1

14 µl of 2.5mM dNTP mix

3.0 µl of Primer 1 (10 µM) (WBR-C5)

30 3.0 µl of Primer 2 (10 µM) (WBR-C3)

1.5 µl of Enzyme (5 units)

63.5 µl water

5 µl DNA

Reaction conditions: 5 cycles ( 93°C, 1 Min; 72°C, 40s; 60°C, 45 sec; 68°C, 2 min) and 25 cycles ( 93°C, 1 min; 72°C, 1 min; 68°C, 2 min).

The amplified product was subcloned into Kpn I and Xho I sites of pcDNA3 vector (Invitrogen). Plasmid DNA was prepared using the method described above.

## EXAMPLE 2

### Functional Assay

Cos-1 cells were transfected as described in Analytical Biochemistry, 218:460-463(1994) with Rat clone 587 GLP-2 receptor, cloned human GLP-2 receptor (pC3/HuGL2R-2), or cloned residue 85 variant human GLP-2 receptor (pC3.1/HuGL2R-MH4), pcDNA3. Rat GLP-1 (7-36) amide was used as a control peptide. Solutions used were as follows:

RSC in RPMI 1640 (49ml RPMI + 1ml FCS + 50ul chloroquine, 100mM);

DEAE/RSC Solution: 18.4 ml RSC + 1.6 ml DEAE/Dextran (10 mg/ml).

The assay procedure entailed the following:

a) 50 mg of either rat clone 587 GLP-2 receptor, or cloned human GLP-2 receptor, or cloned residue 85 variant human GLP-2 receptor was added (as plasmid pcDNA3) to a 50 ml tube containing six mls. of RSC and incubated at 37°C.

b) Six ml of DEAE/RSC solution was added to each tube and incubated at 37°C for 2 min.

c) 1.5 ml of COS-1 cell suspension (5.5 millions cells) was added to each tube and incubated for 1 hr 45 min at 37°C.

d) Following incubation, the sample was spun for 5 minutes at low speed, washed with DMEM/F12 + 10% FBS twice, and the pellet resuspended in 12.5 ml DMEM/F12 + 10% FBS media.

e) One ml of cell suspension (step d) was added to each well of 6 well plates coated with poly- D-lysine (from Collaborative Biomedical), containing 3 ml of media (0.45 million cells/well).

f) Plates were incubated at 37°C for 3 days.

Treatment of Transfected Cos-1 cells with GLP-1/GLP-2 analog was done as follows:  
Solutions : DMEM/F12 (SFM)+ IBMX (3-isobutyl-1-methylxanthin) 0.85mM +0.1%  
5 ascorbic acid and 10 um pargyline (all solutions purchased from Sigma). Media was prepared fresh on day of use.

Assay Procedure: The culture media of each well (transfected 6 well plates, cells) was removed, and the wells were washed once with SFM media. Then 2 ml of SFM +  
10 IBMX media was added to each well and plates were incubated at 37°C for 10 min.

Following incubation, the SFM + IBMX was removed from each well and fresh SFM +  
15 IBMX media containing GLP-1/GLP-2 (GLP-1,7-36,amide from Sigma, [Gly2]hGLP-2 from Allelix) concentration 1, 3, 10 and 30 nM were added to the appropriate wells. Plates incubated at 37°C incubator for 30 min. Following incubation, the media were removed from each well. The wells were washed once with 1ml PBS (Phosphate  
20 Buffered Saline). Each well was then treated with 1ml cold 95% ethanol: 5mM EDTA (2:1) at 4°C for 1 hr. Cells from each well then were scraped and transferred into individual eppendorf tubes. Tubes were centrifuged for 5 min at 4°C, and the supernatants were transferred to new eppendorf tubes and dried in speed vacuum. Following drying, tubes were reconstituted in 100ul of Na- Acetate and kept at 4°C, 25  
20 µl of this solution used for cAMP Assay.

The functional assay was performed as follows: cAMP content for each extract was determined in duplicate by EIA (Enzyme ImmunoAssay) using the Amersham Biotrak cAMP EIA Kit (Amersham 225). Results of the assays, illustrated in Figure 3 and  
25 Figure 8, demonstrate the GLP-2 selectivity exhibited by the cloned rat and human receptors. In a similar functional assay used to users binding to the GLP-1 receptor, the expected specificity for GLP-1 was observed.

### EXAMPLE 3

#### Isolation Of Human GLP-2 Receptor cDNA

#### Medium-stringency hybridization screening of a human hypothalamus cDNA library

One million clones from a  $\lambda$ gt10 cDNA library from human hypothalamus (Clontech; Cat. No. 1172a) were screened by plaque lifts on nitrocellulose filters (Amersham; Cat.RPN137E). The probe was prepared by random primer labelling of a DNA fragment containing the complete coding region of rat GLP-2 receptor. The DNA fragment was isolated from clone 587-C1, which contains the complete coding region from SEQ ID NO: 2.

Pre-hybridization and hybridization were each carried out overnight in a hybridization solution consisting of 50% formamide, 5X SSPE, 5X Denhart's solution, 0.5% SDS and salmon sperm DNA (200 mg/ml). After hybridization the filters were washed under the following conditions (medium stringency):

two times at room temperature in 2X SSPE and 0.01% SDS.

two times at 42°C in 2X SSPE and 0.01% SDS.

two times at 42°C in 0.2X SSPE and 0.01% SDS.

The filters were autoradiographed and agar plugs, each containing numerous bacteriophage plaques, were picked from regions on the plates corresponding to positive signals on the filter. From one million cDNA clones sampled in the first round screen, there were identified two positive clones (HHP6-1 and HHP13). On secondary screening only HHP13 turned out positive. Several positive plaques (HHS13) from the HHP13 plate were pooled and taken for tertiary screening. Three single positive plaques from this round of screening were picked (HHT13-1, HHT13-2, HHT13-3).

PCR amplification was then used for partial sequencing of the positive clones. On a lawn of bacterial cells (E.coli C600Hfl), 10  $\mu$ l of phage suspension from each clone was applied at marked spots. After 5 hr incubation at 37°C, the phage plaques were clearly visible and covered  $\sim 1$  cm<sup>2</sup>. A portion of each plaque was transferred to 200  $\mu$ l of water. The samples were incubated in a boiling water bath for 5 min and centrifuged at room temperature for 10 min. One millilitre of sample was used for PCR amplification with two sets of degenerate primers:

M2FS[5'-TTTTTCTAGAASRTSATSTACACNGTSGGCTAC-3'] (SEQ ID NO: 3) and

M7RS[5'-TTTTCTCGAGCCARCARCCASSWRTARTTGGC-3'] (SEQ ID NO: 4); or  
C4-4 [5'-AACTACATCCACMKGMAYCTGTTYVYGTCBTTCATSCT-3'] (SEQ ID  
NO: 5) and

C9-2R [5'-TCYRNCTGSACCTCMYYRTTGASRAARCAGTA-3'] (SEQ ID NO: 8).

The Expand™ PCR system from Boehringer Mannheim (Cat. 1681-842) was used under  
the following conditions:

5 µl of 10x Expand™ Buffer 3

7 µl of 2.5mM dNTP mix

1.5 µl of primer M2FS or C4-4

1.5 µl of primer M7RS (with M2FS) or C9-2R (with C4-4)

0.75 µl of Expand PCR enzyme (1 unit)

33.25 µl water and

1 µl DNA.

Reaction conditions were: 32 cycles at 93°C, 1 min; cycles 50 °C, 1 min; cycles 45 °C, 1  
min; cycles 68°C, 2 min.

M2F/S and M7R/S amplified a DNA fragment of about 300 bp and C4-4 and C92-R  
amplified a DNA fragment of about 700 bp. The PCR products were purified using the  
QIAGEN QIAquick PCR purification kit (Cat. 28104) and eluted in 50 µl 10 mM Tris,  
pH 8.0. Sequence analysis of the products revealed no differences between the  
templates, as expected from the fact that they represent multiple copies of a single cDNA  
clone (HHT13).

A number of factors indicate that this clone contains coding sequence of the human  
GLP-2 receptor. One factor is the degree of sequence similarity. The glucagon receptor  
cDNA can be used to predict the expected degree of sequence conservation found  
between rat and human receptors. At the nucleotide level, there is 82.6% identity within  
the coding regions of the rat and human glucagon receptors. At the amino acid level,  
there is 80.9% identity and 89.1% amino acid similarity between the glucagon receptors  
of the two species.

In the case of the human GLP-2 receptor cloned herein, the sequence of the partial human GLP-2 receptor cDNA (HHT13) is highly homologous to rat GLP-2 receptor cDNA at both the nucleotide and amino acid level. SEQ ID NO: 9 shows 87.1% identity with the rat GLP-2 receptor cDNA sequence. The predicted amino acid sequence of this cDNA region has 87.4% identity and 93.2% similarity with the predicted amino acid sequence of the rat GLP-2 receptor. The total predicted length of the rat receptor preprotein is 550 amino acids, suggesting about 44% of the coding region of the human receptor had been identified.

Further evidence supporting this conclusion comes from a comparison of the partial human GLP-2 receptor amino acid sequence with the rat GLP-2 receptor and the 3 next closest family members, shown below:

Receptor Sequence (amino acid)	Percent Identity with HHT13	Percent Similarity
GLP-2 receptor (rat)	87.4	93.2
GLP-1 receptor (rat)	50.0	74.1
Glucagon receptor (rat)	51.4	73.9
GIP receptor (rat)	50.7	70.3

These comparisons, together with the benchmark provided by sequence similarities between the rat and human glucagon receptors, provide definitive evidence that the cDNA designated HHT13 represents a fragment of the human counterpart of the rat GLP-2 receptor.

The full amino acid sequence of the human GLP-2 receptor can be obtained by first determining the sequence of the complete cDNA inserts in HHT13-1, HHT13-2 and HHT13-3. By using degenerate primers for PCR amplification and subsequent sequencing, we obtained sequence from only part of each insert. It is possible that these identical clones contain an insert which spans the complete coding sequence of the human GLP-2 receptor preprotein. To determine the complete sequence of the cDNA insert, the clones are grown in large quantity to prepare approximately 20 mg of each



equivalent clone. The complete cDNA insert is excised by restriction with Eco RI, and subcloned into pcDNA3 (Invitrogen). Alternatively, two primers from vector sequence flanking the insert are used to amplify the complete cDNA insert using the Expand™ PCR system from Boehringer Mannheim (Cat. 1681-842). The amplified cDNA is cut with appropriate restriction enzymes and is subcloned into pcDNA3 (Invitrogen).

If a complete coding sequence is not present in the HHT13 clones, cDNA libraries are screened for additional clones to complete the coding region of human GLP-2 receptor cDNA. Preferably human cDNA libraries (from Stratagene or Clontech) representing the following tissues are used for screening: Human hypothalamus; Human fetal brain; Human duodenum and jejunum; Human stomach; and Human fetal intestine.

Two PCR primers are designed from the sequence of human GLP-2 receptor cDNA already determined. These primers are designed such that they could not amplify any related gene family members other than the GLP-2 receptor cDNA itself. A dilution of the cDNA library stock is used to make library sub-pools such that 50,000 clones are represented in each pool. PCR is conducted with the GLP-2 receptor-specific primers to diagnose pools containing a GLP-2 receptor cDNA clone, using the Expand™ PCR system from Boehringer Mannheim (Cat. 1681-842) under the following conditions:

2 µl of 10x Expand™ Buffer 1

2.8 µl of 2.5mM dNTP mix

0.6 µl of primer P1

0.6 µl of primer P2

0.3 µl of Expand PCR enzyme (1 unit)

12.7 µl water

1 µl of library pool containing 50,000 clones

Reaction conditions: 32 cycles at 93°C, 40 sec; 50-58°C, 40 sec; 68°C, 40 sec.

Sequence is then obtained from the complete GLP-2 receptor cDNA insert from a positive pool. By using one specific primer and one primer based on vector sequence close to the cloning site, the GLP-2 receptor cDNA insert is directly amplified from

clonally impure clone pools, using the Expand™ PCR system from Boehringer Mannheim (Catalogue no. 1681-842) most suitably under the following conditions:

2 µl of 10x Expand™ Buffer 1

2.8 µl of 2.5mM dNTP mix

5 0.6 µl of Primer 1

0.6 µl of Primer 2

0.3 µl of Enzyme (1 unit)

12.7 µl water

1 µl of library pool stock

10 Reaction conditions: 32 cycles at 93°C, 45 sec; 50°C, 45 sec; 68°C, 1 min.

The reaction is run on a preparative agarose gel, and the most prominent band is purified and sequenced. Based on the amplified sequence obtained, additional primers are designed to obtain sequence and clones of complete coding region and clone the complete cDNA

15

5' RACE and 3' RACE are used to obtain complete coding sequence of the human GLP-2 receptor cDNA. Rapid Amplification of cDNA Ends (RACE) is a procedure routinely used for amplification of DNA sequences from first cDNA strand (easily prepared from mRNA) template between a defined internal site and either 3' or the 5' end of the

20

mRNA. Total or mRNA from different human tissues are commercially available from Clontech. The 3' RACE System (Gibco-BRL Life Technologies; Cat. 18373-019) and 5' RACE System (Cat. 18374-058) kits are used. The manuals of these two products provide detailed protocols. In brief, protocols are as described below.

25

For the 3' RACE procedure, first strand cDNA synthesis is initiated at the poly (A) tail of mRNA using the adapter primer (provided with system) incorporating a unique sequence for universal PCR amplification of the RACE products. After synthesis of the first strand cDNA from this primer, the original mRNA template is destroyed with RNase H. Amplification is then performed using two primers: one is a gene-specific

30

primer (which will be designed from the available partial cDNA sequence of HHT13);

the other is the universal amplification primer provided with the kit. The amplified product is subcloned into a plasmid vector for sequencing.

For the 5' RACE System, the first strand cDNA is synthesized from mRNA using a gene-specific primer (which is based on the available partial cDNA sequence of HHT13) and SuperScript II reverse transcriptase. The original mRNA template is removed by treatment with RNase H. Unincorporated dNTPs, primer, and proteins are separated from cDNA using spin cartridge. A homopolymeric dCTP tail is then added to the 3'-end of the first strand cDNA using TdT enzyme and dCTP nucleotides. PCR amplification is performed using two primers: one is a nested, gene-specific primer designed from the available partial DNA sequence of HHT13; and the other is an "anchor primer" provided with the system. Both primers incorporate restriction sites for subcloning into plasmids and subsequent sequencing.

Sub-cloning of HHT13  $\lambda$ gt10 clones into pcDNA3, their sequencing and expression

A. Amplification of cDNA inserts with  $\lambda$ gt10 primers.

On a lawn of bacterial cells (E.coli C600Hfl), 10  $\mu$ l of phage resuspension from each clone was placed at marked spots. After 5 hr incubation at 37°C, the phage plaques were clearly visible. The surface of each plaque was transferred to 200  $\mu$ l of water. The samples were kept in boiling water bath for 5 minutes and centrifuged at room temperature for 10 minutes. 1  $\mu$ l of sample was used to amplify with a set of  $\lambda$ gt10 primers.

GT10-5KXb [5'-GGGTAGTCGGTACCTCTAGAGCAAGTTCAGCC-3']

vs

GT10-3BXh [5'-ATAACAGAGGATCCTCGAGTATTTCTTCCAG-3']

The Expand™ PCR system from Boehringer Mannheim (Cat. 1681-842) was used under the following conditions:

5  $\mu$ l of 10x Expand™ Buffer 3

7  $\mu$ l of 2.5mM dNTP mix

1.5 µl of primer GT10-5KXb  
1.5 µl of primer GT10-3BXh  
0.75 µl of Expand PCR enzyme (1 unit)  
33.25 µl water  
5 1 µl DNA

Reaction conditions: 5 cycles of 93°C-40 sec; 50°C- 1 min; 68°C-2 min and 30 cycles of 93°C-40 sec; Ramp to 68°C- 1 min; 68°C-2 min.

10 An amplified DNA fragment of about 2200 bp long was seen on the agarose gel from all three clones. The PCR product were purified using the QIAGEN's QIAquick PCR purification kit (Cat. no. 28104) and eluted in 50 µl 10 mM Tris, pH 8.0. The templates were sequenced.

#### 15 B. Subcloning into pcDNA3 vector.

The amplified and purified DNA from the three clones was restricted with Kpn I and Xho I and subcloned into pcDNA3 restricted with similar restriction enzymes. The plasmids were named pHHT13-1, pHHT13-2, and pHHT13-3. Plasmids DNAs were prepared using either crude method (alkaline treatment, bacterial DNA precipitation with 3 M KOAc, isopropanol precipitation followed by RNase treatment and second round of  
20 isopropanol precipitation) or plasmid DNA kits from Qiagen Inc. The templates prepared using Qiagen's kits were sequenced.

#### C. Functional Assay

25 Transfections were carried out with each clone, using the rat GLP-2R, 587 clone as a positive control for cAMP response to GLP-2 peptide. Methods for transfection, cell culture and cAMP assay were identical to those described for the functional assay of rat, 587 clone. Results showed that although the positive control gave good cAMP response in COS cells, none of the HHT13 clones gave any cAMP response. As confirmed by sequencing which showed a frame-shift mutation, the functional data suggested that no  
30 functional GLP-2R protein was expressed from these cDNA clones.

D. Comparison of DNA sequences between rat GLP-2R and HHT13 subclones.

The comparison showed a 2 bp deletion at a position corresponding to nucleotides 389-390 of the rat GLP-2R cDNA, resulting in the loss of nucleotides 374-375 of the human GLP-2R cDNA sequence presented herein.

5 PCR was used to incorporate two bp of the rat GLP-2R DNA sequence into HHT13-1 DNA at the site of the 2 bp frame-shift deletion identified relative to the rat GLP-2R coding sequence. The following primers were designed from HHT13 DNA sequence to insert two bp :

10 HWBR/2BPI-475F

[5'-ACAGGCATGTCTGGAAGACTTACTCAAGGAACCTTCTGGCAT-3']

HWBR/2BPI-506R

[5'-ATGCCAGAAGGTTCTTGAGTAAGTCTTCCAGACATGCCTGT-3']

HWBR-F7 [5'-TTCCTCTGTGGTACCAAGAGGAATGC-3']

15 and HWBR-1910R:

[5'-GGTGGACTCGAGGTACCGATCTCACTCTCTTCCAGAATC-3']

PCR 1: One ng of pHHT13-1 DNA was used as template to do two PCRs with primers, HWBR-F7 vs HWBR/2BPI-506R and HWBR/2BPI-475F vs HWBR-1910R.

20 The Expand™ PCR system from Boehringer Mannheim (Cat. 1681-842) was used under the following conditions:

5 µl of 10x Expand™ Buffer 1

7 µl of 2.5mM dNTP mix

1.5 µl of primer HWBR-F7 or HWBR/2BPI-475F

25 1.5 µl of primer HWBR/2BPI-506R or HWBR-1910R

0.75 µl of Expand PCR enzyme (1 unit)

33.25 µl water and

1 µl DNA.

30 Reaction conditions: 10 cycles of 92°C-40 sec; 48°C- 1 min; 68°C-3 min and 30 cycles of 92°C-40 sec; 55°C- 40 sec; 68°C-2 min.

The primers HWBR-F7 and HWBR/2BPI-506R amplified a DNA fragment of 400 bp and HWBR/2BPI-475F and HWBR-1910R amplified a DNA fragment of approximately 1.4 kb on an agarose gel. The two bands were cut out of the agarose gel and purified with Qiaquick gel extraction kit from Qiagen Inc. (Cat no. 28706) and the DNAs were eluted in 50 µl of 10 mM Tris (pH 8.5).

PCR 2 (Extension without primers): Approximately 75 ng of two amplified product from above PCR 1 were mixed and then recombined without primers by extending under the following conditions:

2 µl of 10x Expand™ Buffer 1

2.8 µl of 2.5mM dNTP mix

0.3 µl of Expand PCR enzyme (1 unit)

8.9 µl water

6 µl of combined PCR 1 products

Reaction conditions: 15 cycles of 92°C-1 min; 60°C- 5 min; 68°C-3 min.

PCR 3: 1 µl of amplified mix from PCR 2 was used as template to amplify with HWBR-F7 and HWBR- 1910R primers using the following conditions:

10 µl of 10x Expand™ Buffer 1

14 µl of 2.5mM dNTP mix

3.0 µl of primer HWBR-F7 or HWBR/2BPI-475F

3.0 µl of primer HWBR/2BPI-506R or HWBR-1910R

1.5 µl of Expand PCR enzyme (1 unit)

67.5 µl water and

1 µl DNA.

Reaction conditions: 30 cycles of 92°C-1 min; 60°C- 1 min; 68°C-2 min.

A DNA fragment of approximately 1.7 kb was amplified as seen on an agarose gel. The PCR product was purified using the QIAGEN's QIAquick PCR purification kit (cat. no. 28104) and eluted in 50 µl of 10 mM Tris, pH 8.0. The purified product was restricted with Kpn I and subcloned into Kpn I-restricted pcDNA3.1(-)/Myc-His A (Invitrogen,

Cat. no. V855-20). One clone, named pc3.1/HuGL2R/MH6 (pHuMH6), had the 1.7 kb insert in correct orientation as checked by PCR using vector vs. insert primers.

Functional assay:

5 This hybrid clone was compared to rat GLP-2R using the assay described in Example 2. Results showed that the 2 bp "GA" replacement into the putative deletion site yielded a clone encoding a functional GLP-2R protein, as shown by the cAMP response to GLP-2 treatment.

10 **EXAMPLE 6**

**Isolation of the Full-Length Human GLP-2 Receptor cDNA**

Twenty thousand clones from  $\lambda$ gt10 cDNA Library from Human Stomach (Clontech; Cat. HL3017a) were plated on each of 100 agar 150 mm plates. SM buffer (0.1 M NaCl, 10 mM  $Mg_2SO_4$ , 35 mM Tris, pH-7.5, 0.01% gelatin) was added to each plate to obtain 15 100 phage lysates each containing 20,000 (20K) pooled clones. The first fifty 20K phage lysates (20K pools) were screened by PCR using two primers designed from HHT13 DNA sequence. The template DNA from each pool was prepared by boiling phage lysate for 10 minutes and centrifuging for 10 minutes.

HWBR-113F [5'-GTGGAGAGGATTTGTGCAAACATTTC-3']

20 HWBR-578R [5'-AGAGACATTTCCAGGAGAAGAATGAG-3']

1  $\mu$ l of each 20K pool DNA was diagnosed by PCR with HWBR-113F and HWBR-578R primers using the following conditions:

2  $\mu$ l of 10x Expand<sup>TM</sup> Buffer 1

2.8  $\mu$ l of 2.5mM dNTP mix

25 0.6  $\mu$ l of primer HWBR-113F

0.6  $\mu$ l of primer HWBR-578R

0.3  $\mu$ l of Expand PCR enzyme (1 unit)

12.7  $\mu$ l water

1  $\mu$ l 20K pool DNA

30 Reaction conditions: 35 cycles of 92°C- 40s.; 60°C- 40s.; 68°C-1 min.

A DNA fragment of approximately 450 bp was seen in amplification of templates from two pools (HST 19 and HST 38).

B. Screening of clones from two positive pools: HST 19 and HST 38.

40,000 clones plated from each of two positive 20K pools were screened by plaque lifts on nitrocellulose filters (Amersham; Cat.RPN137E). The probe was prepared by random primer labelling a DNA fragment from pHHT13-1.

1. The filters were pre-hybridized and hybridized at 42°C overnight. Hybridization solution consisted of 50% formamide, 5X SSPE, 5X Denhart's solution, 0.5% SDS and salmon sperm DNA (200 mg/ml).

2. After hybridization the filters were washed under the following conditions:

- two times at room temperature in 2X SSPE and 0.01% SDS;
- two times at 42°C in 2X SSPE and 0.01% SDS; and
- two times at 50°C in 0.1X SSPE and 0.01% SDS.

3. The filters were autoradiographed and the regions on the plates matching to positive signals were isolated. One positive clone (HST 38-4-30) was isolated from HST 38 pool. 450 bp DNA fragment was amplified from the positive clone by using primers HWBR-113F and HWBR-578R and sequenced. The sequence clearly showed that the plasmid contain 2 bp (4G) at position 373-374 of HHT13 DNA sequence.

The complete insert of clone HST 38-4-30 was amplified using  $\lambda$ gt10 primers as described in Example 1. PCR amplified a DNA fragment of approximately 1.4 kb. The amplified DNA was purified and sequenced.

#### **EXAMPLE 7**

##### **Reconstruction of a clone of full-length functional human GLP-2R cDNA and functional assay**



A 700 bp fragment obtained by Kpn I and Pvu II restriction digest of the amplified DNA from clone HST 38-4-30, and 1.4 kb DNA fragment from Xho I and Pvu II restricted pHHT13-1 DNA were subcloned into Kpn I and Xho I restricted pcDNA3 in a three-way ligation. The new plasmid construct was called pc3/HuGL2R-2. In this manner the full length sequence of the human GLP-2 receptor was obtained.

#### Functional Assay:

The new clone was compared to the rat GLP-2R clone 587 as described previously above. Results showed that the clone encoded a functional human GLP-2R protein, which led to cAMP production in COS cells in response to GLP-2 treatment (Figure 8).

### **EXAMPLE 8 Antibodies directed to the GLP-2 receptor**

#### **1. Antipeptide antibodies**

Antipeptide antibodies were raised in rabbits against an N-terminal peptide (QTRENTTDIWQDESE), a C-terminal peptide (SEGDGSETLQKLR) and extracellular loop 1 (SHNSUSKRPDDESG) of the rat GLP-2 receptor.

Immunocytochemical analysis of serum produced as above confirmed that serum contained antibodies directed to the GLP-2 receptor which do not cross react with the rat GLP-1 receptor.

#### **2. Antibodies to a GLP-2 receptor raised against a fusion protein**

Polynucleotide encoding the C-terminal region of rat GLP-2 receptor (amino acids 444-550) was spliced to the C-terminus of glutathione S-transferase (GST) in pGEX-2T and expressed in E.coli strain SURE1. Protein was purified using affinity chromatography using the above GLP-2 C-terminal fragment fused to the C-terminus of maltose binding protein. Protease degradation was minimized by using a cocktail of protease inhibitors (Boeringer Mannheim).

Antibodies were raised generally according to the method disclosed in Antibodies: A laboratory manual, Harlow and Lane, Cold Spring Harbor Laboratory, 1988. Briefly, the

5 GLP-2-GST fusion protein was used to raise antibodies in rabbits as follows. Initial injection was with 100 µg of fusion protein in complete Freund's adjuvant at multiple sites, intramuscularly and subcutaneously. Booster injections were made at multiple sites intramuscularly with 100 µg of fusion protein in incomplete Freund's adjuvant at days 14, 21, 42 and 56.

Antisera was affinity purified using a GLP-2-MBP fusion protein affinity column. Immunocytochemical analysis confirmed that these antibodies specifically recognize the GLP-2 receptor.

#### EQUIVALENTS

10 The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. Indeed, various modifications of the above-described makes for carrying out the invention which are obvious to those skilled in the field of biochemistry, molecular biology or related fields are intended to be within the scope of the following claims.

What is claimed is:

1. An isolated polynucleotide encoding a GLP-2 receptor.
- 5 2. The isolated polynucleotide according to claim 1, which codes for the human GLP-2 receptor having the amino acid sequence of amino acids 67-553 of SEQ ID NO:12
3. The isolated polynucleotide according to claim 2, comprising nucleotides 318-1780 of SEQ ID NO:11
- 10 4. The isolated polynucleotide according to claim 1, which codes for the rat GLP-2 receptor having the amino acid sequence of amino acid 67-550 of SEQ ID NO:2
5. The isolated polynucleotide according to claim 4, having nucleotides 335-1789 of SEQ ID NO: 1.
- 15 6. An isolated polynucleotide which hybridizes under conditions of high stringency with a polynucleotide defined in claim 3 or claim 5.
- 20 7. An oligonucleotide comprising at least 15 nucleotides and having a polynucleotide sequence that hybridizes selectively to a polynucleotide according to any of claims 1-6.
8. An oligonucleotide according to claim 7, wherein the nucleotide sequence of the oligonucleotide corresponds identically with a region of the polynucleotide defined in claim 3 or claim 5.
- 25 9. An oligonucleotide according to claim 7, wherein the nucleotide sequence of the oligonucleotide corresponds identically with a region common to the polynucleotides defined in claim 3 and claim 5.

30

10. In labelled form, a polynucleotide selected from a polynucleotide as defined in any of claims 1-6, and an oligonucleotide as defined in any of claims 7-9.

5 11. A recombinant polynucleotide comprising a GLP-2 receptor-encoding polynucleotide as defined in any of claims 1- 6, and expression controlling elements linked operably therewith to drive expression thereof.

12. A cell that has been genetically engineered by the incorporation expressibly therein of a polynucleotide according to any of claims 1-6.

10 13. The cell according to claim 11, which is a mammalian cell.

14. An antibody which selectively binds to a GLP-2 receptor.

15 15. A recombinant GLP-2 receptor.

16. The GLP-2 receptor of claim 15, comprising amino acids 67-553 of SEQ ID NO:2.

17. The GLP-2 receptor of claim 15, comprising amino acids 67-550 of SEQ ID NO: 12

20 18. A method for identifying GLP-2 receptor ligands, which comprises the steps of:

25 (1) incubating a candidate ligand with a cell as defined in claim 12 or with a membrane preparation obtained therefrom; and then

(2) determining whether binding between the GLP 2 receptor and the candidate ligand has occurred.

30 19. A GLP-2 receptor ligand identified by the method of claim 18.

1/13

1 AAGCTTCGCG GCCTCTGCAG AKGAGTTGTG CAAACACTTC CTCTCTGGAC  
51 AAGGAGGAAT GCAGGAGGCC ACCGCCTGCA GTACATCTTG GAGTGTGGGA  
101 GGGATGTGCC TGCACCTGTG AACGGGCGCC AGGAGAATGA GGCCCAACC  
151 AAGCCCGCA GTGCCAGTA GATGCAGAGA GGCACCCGTG CCCCAGTGTA  
201 GGGCACAGCC AGTGGGCATC CCTGAGGCC AGGGGCCGT TCCTCTCCAC  
251 TCCCAACAGA TCGTCTGCT GTGGGGCCCT GGGAGGCCCT TCCTCGCCCT  
301 GCTTCTGCTG GTTTCCATCA AGCAAGTTAC AGGATCGCTC CTCAAGGAGA  
351 CAACTCAGAA GTGGGCTAAT TATAAGGAGA AGTGTCTGGA AGACTTGCAC  
401 AATAGACTTT CTGGCATATT TTGTAATGGG ACATTTGATC GGTATGTGTG  
451 CTGGCCTCAT TCTTATCCTG GAAATGTCTC TGTTCCTGT CTTTCACT  
501 TACCTTGGTG GAATGCAGAG AGCCCAGGAA GGCCTACAG AACTGCTTG  
551 GCTCAGGGGA CTTGGCAGAC GCGAGAGAAC ACCACAGATA TTTGGCAGGA  
601 TGAATCAGAA TGCTCAGAGA ACCACAGCTT CAGACAAAAC GTGGATCACT  
651 ACGCCTTGCT ATACACCTTG CAGCTGATGT AACTGTGGG CTACTCCGTG  
701 TCTCTCATCT CCCTCTTCTT GGCTCTTACA CTCTTCTTGT TCCTTCGAAA  
751 ACTGCATTGC ACACGCAATT ACATCCACAT GAACCTGTTC GCTTCGTTCA  
801 TCCTGAAAGT TCTGGCTGTC CTGGTGAAGG ACATGGTCTC CCACAACCTCT  
851 TACTCCAAGA GGCCCGATGA TGAGAGTGA TGGATGTCAT ATCTGTCAGA  
901 GACATCCGTC TCCTGTGCT CCGTCCAGGT CCTCCTGCAC TACTTTGTGG  
951 GCACCAATCA CTTGTGGCTG CTGGTTGAAG GACTTTACCT CCACACTCTG  
1001 CTGGAGCCCA CAGTGTTC TGAAGGCGG CTGTGGCCCA AGTACCTGGT  
1051 GGTGGGTTGG GCCTTCCCCA TGCTGTTTGT TATTCCCTGG GGTTCGCCC  
1101 GTGCACACCT GGAGAACACA CCGTGCTGGG CCACAAATGG GAACCTGAAA  
1151 ATCTGGTGGA TCATCAGAGG ACCCATGCTG CTTTGTGTAA CAGTTAATTT  
1201 CTTTCTTCTC CTCAAGATTC TCAAGCTTCT CATTTCTAAG CTCAAAGCTC  
1251 ATCAGATGTG CTTTCTTCTC TCAAAATACA GATTGGCGAA ATCAACGTTG  
1301 CTCCTCATTC CTTTGTGGG GTTTCATGAG GTCCTCTTCA CTTTCTTCCC  
1351 CGACGACCAA GTTCAAGGAT TTTCAAAACG TATTGACTC TTCATCCAGC  
1401 TGACACTGAG CTCTGTCCAC GGATTTCTGG TGGCCTTGCA GTATGGCTTT  
1451 GCCAATGGAG AGGTGAAGGC AGAGCTGCGA AAGTCATGGG GCCGCTTCTT  
1501 ATTAGCCCGC CACTGGGGCT GCAGAACCCTG TGTCTGGGG AAGAATTTCC  
1551 GGTTCTTGGG GAAGTGTTC AAGAAGCTGT CGGAGGGAGA TGGCTCTGAG  
1601 AACTCCAGA AGCTGCGGTT CTCCACATGC AGCTCACACC TGGCCTCTGA  
1651 GACCTGGGA GACGTTGGG TACAGCCTCA CAGGGGCCGT GGAGCTTGGC  
1701 CCCGGGAAG CAGCCTGTCT GAGAGCAGTG AGGGAGACTT CACCCTGGCC  
1751 AATACGATGG AGGAGATTCT GGAAGAGAGT GAGATCTAAG GCAGGTCCA

FIG.1A

2/13

1801 TCACCGCAGC TTGGCCACTG ARGAmCCAAC CcTARGAAGG ATKTTGCCGA  
1851 RCCCARGGTC CTCCTCTTCC TATGTWCtAT MCCCATTTTG ATGTGAAGTC  
1901 TCTCCTGGGT GAMCAASCTC TGTACCAACS ARTCTCAgTC CCTCTTGCCC  
1951 TTGTCAcCCT ACTACCCCTC CCCCATCαCα CATgTTTTCC AGAATKTCCG  
2001 TtGGTTTGGG GGGGGGGGTC TTGCCCTAAA TTCAAGTsGA GTGGARCCCA  
2051 CCATGAAGAA AARTCATTTA TTAAATAGAR TCCGGTTAGG ATCTCCTTCC  
2101 CGTTCATGGT GCATGGCCTC CTTCCAAGGG ATGGGAGTCG GSTGCACTGG  
2151 AACCACACAG GAAAcTTTGA AGTATCCAGT TCTAGGGAAT TATAGCCAAT  
2201 ATTCTGAGAG AGCAAGTCTG AGATGAGAKC CGAGAATWGC AAGTGTWGGA  
2251 CAWGCATTCA AGGAAACTCC TCACCTTTgG GCGAAACCTA tGGCAGGATc  
2301 GGCAtGGAGC AGCTATTMTG CAAYGGCCGC TCACCTGGGA CATACTACTC  
2351 TCCTTGGGCA GGATGTGACC CCATGTKGTC CCCAGACTC CTCTCCTCCT  
2401 TGCTTSTSTT CYTTTCCYGT CAAGTCTCAC CTCCCTTTCT ACATCTCAGT  
2451 TCWGTTTGGT GTYGACAGAA GYYTGAATGT CACAATACTG CATGTGTTAG  
2501 TTTCTGTCGT CATTGCTGTG TCCAAATACC TGACCAGGAC CAATTTAAGC  
2551 GAGGAAGTGC TACATGGGCG GCCGC

FIG.1B

3/13

1 MRPQSPAVP SRCREAPVPR VRAQPVGIPE AQGPVPLHSQ QMRLLWGPGR  
51 PFLALLLLVS IKQVTGSLLK ETTQKWANYK EKCLEDLHNR LSGIFCNGTF  
101 DRYVCWPHSY PGNVSVPCPS YLPWWNAESP GRAYRHCLAQ GTWQTRENTT  
151 DIWQDESECS ENHSFRQNV D HYALLYTLQL MYTVGYSVSL ISLFLALTLF  
201 LFLRKLHCTR NYIHMLFAS FILKVLAVLV KDMVSHNSYS KRPDDESGWM  
251 SYLSETSVSC RSVQVLLHYF VGTNHLWLLV EGLYLHTLLE PTVFPERRLW  
301 PKYLVVGWAF PMLFVIPWGF ARAHLENTRC WATNGNLKIW WIIRGPMLLC  
351 VTVNFFIFLK ILKLLISKLK AHQMCFRDYK YRLAKSTLLL IPLLGVHEVL  
401 FTFFPDDQVQ GFSKRIRLFI QLTLSVHGF LVALQYGFAN GEVKAELRKS  
451 WGRFLLARHW GCRTCVLGKN FRFLGKCSKK LSEG DGSETL QKLRFCSTCS  
501 HLASETLGDV GVQPHRGRGA WPRGSSLSSES SEGDFTLANT MEEILEESEI

FIG.2

4/13




FIG.3



5/13

```
1 TCCTTCTCTC TTATCTCCCT CTCCTGGCT CTGACCCTCC TCTTGTTTCT
51 TCGAAAACTC CACTGCACGC GCAACTACAT CCACATGAAC TTGTTTGCTT
101 CTTTCATCCT GAGAACCCTG GCTGTACTGG TGAAGGACGT CGTCTTCTAC
151 AACTCTTACT CCAAGAGGCC TGACAATGAG AATGGGTGGA TGTCTTACCT
201 GTCAGAGATG TCCACCTCCT GCCGCTCAGT CCAGGTTCTC TTGCATTACT
251 TTGTGGGTGC CAATTACTTA TGGCTGCTGG TTGAAGGCCT CTACCTCCAC
301 ACGCTGCTGG AGCCACAGT GCTTCCTGAG AGGCGGCTGT GGCCCARATA
351 CCTGCTGTTG GGTGGGCCT TCCCTGTGCT ATTTGTTGTA CCCTGGGGTT
401 TCGCCCGTGC ACACCTGGAR AACACAGGGT GCTGGACAAC AAATGGGAAT
451 AAGAAAATCT GGTGGATCAT CCGAGGACCC ATGATGCTCT GTGTAACAGT
501 CAATTTCTTC ATCTTCCTGA AAATTCTCAA GCTTGTCATT TCTAAGCTCA
551 AAGCTCATCA AATGTGCTTC AGAGATTATA AATACAGATT GGCAAAATCA
601 ACACTGGTCC TCATTCCTTT ATTGGGCGTT CATGAGATCC TCTTCTCTTT
651 CATCACTGAT GATCAAG
```

FIG.4

6/13

S F S L I S L F L A L T L L L F L R K L  
TCCTTCTCTCTTATCTCCCTCTTCTGGCTCTCACCTCCTCTTGTTCGAAAACCTG  
1 -----+-----+-----+-----+-----+----- 80

H C T R N Y I K M N L F A S F I L R T L  
CACTGCACGCGCAACTACATCCACATGAACCTGTTTGCTTCTTTCATCCTGAGAACCCTG  
61 -----+-----+-----+-----+-----+----- 120

A V L V K D V V F Y N S Y S K R P D N E  
GCTGTACTGGTGAAGGACGTCGTCTTCTACAACCTTACTCCAAGAGGCCTGACAATGAG  
121 -----+-----+-----+-----+-----+----- 180

N G W M S Y L S E M S T S C A S V Q V L  
AATGGGTGGATGTCCTACCTGTCAGAGATGTCCACCTCCTGCCGCTGAGTCCAGGTTGTG  
181 -----+-----+-----+-----+-----+----- 240

L H Y F V G A N Y L W L L V E G L Y L H  
TTGCATTACTTTGTGGGTGCCAATTACTTATGGCTGCTGGTGAAGGCCTCTACCTCCAC  
211 -----+-----+-----+-----+-----+----- 300

T L L E P T V L P E R R L W P K Y L L L  
ACGCTGCTGGAGCCACAGTGCTTCCTGAGAGGGGGCTGTGGCCCA?ATACCTGCTGTTG  
301 -----+-----+-----+-----+-----+----- 360

G W A F F V L F V V P W G F A R A H L E  
GGTTGGGCCTTCCCTGTGCTATTTGTGTACCTGGGGTTTCGCCCCTGCACACCTGGAR  
361 -----+-----+-----+-----+-----+----- 420

N T G C W T F N G N K K I W W I I R G P  
AACACAGGGTGCTGGACAACAAATGGGAATAAGAAAATCTGGTGGATCATCCGAGGACCC  
421 -----+-----+-----+-----+-----+----- 480

M M L C V T V N F F I F L K I L K L L I  
ATGATGCTCTGTGTAACAGTCAATTTCTTCATCTTCCTGAAAATTCTCAAGCTTCTCATT  
481 -----+-----+-----+-----+-----+----- 540

S K L K A H Q M C F R D Y K Y R L A K S  
TCTAAGCTCAAAGCTCATCAAATGTGCTTCAGAGATTATAAATACAGATTGGCAAAATCA  
541 -----+-----+-----+-----+-----+----- 500

T L V L I P L L G V H E I L F S F I F D  
ACACTGGTCTCATTCCTTTATTGGCGTTTCATGAGATCCTCTTCTCTTTCATCACTGAT  
501 -----+-----+-----+-----+-----+----- 560

D Q  
GATCAAG  
661 ----- 667

FIG.5

7/13

1	TGGAGAGGATTTGTGCAAACATTTCTTCTGTGGACCAAGAGGAATGCAAGAGGAGGCTGC	60
61	CTGCGGTGCATCTTGGACGGCTAGAGAGATGTACCCCTACTTGTGAAGGTGCACGAGGAA	120
121	M K L G S S R A G P G R G S A G L L P G GATGAAGCTGGGATCGAGCAGGGCAGGGCCTGGGAGAGGAAGCGCGGGACTCCTGCCTGG	180
181	V H E L P M G I P A P W G T S P L S F H CGTCCACGAGCTGCCCATGGGCATCCCTGCCCCCTGGGGGACCAGTCCTCTCTCCTTCCA	240
241	R K C S L W A P G R P F L T L V L L V S CAGGAAGTGCTCTCTCTGGGCCCCCTGGGAGGCCCTTCTCACTCTGGTCCTGCTGGTTTC	300
301	I K Q V T G S L L E E T T R K W A Q Y K CATCAAGCAAGTTACAGGATCCCTCGTTGAGGAAACGACTCGGAAGTGGGCTCAGTACAA	360
361	Q A C L R D L L K E P S G I F C N G T F ACAGGCATGTCTGAGAGACTTACTCAAGGAACCTTCTGGCATATTTGTACGGGACATT	420
421	D Q Y V C W P H S S P G N V S V P C P S TGATCAGTACGTGTGTTGGCCTCATTCTCTCCTGGAAATGTCTCTGTACCCTGCCCTTC	480
480	Y L P W W S E E S S G R A Y R H C L A Q ATACTTACCTTGGTGGAGTGAAGAGAGCTCAGGAAGGGCCTACAGACACTGCTTGGCTCA	540
541	G T W Q T I E N A T D I W Q D D S E C S GGGGACTTGGCAGACGATAGAGAACGCCACGGATATTTGGCAGGATGACTCCGAATGCTC	600
601	E N H S F K Q N V D R Y A L L S T L Q L CGAGAACCACAGCTTCAAGCAAAACGTGGACCGTTATGCCCTGCTGTCAACCTTGCAGCT	660

FIG.6A

661	M Y T V G Y S F S L I S L F L A L T L L GATGTACACCGTGGGATACTCCTTCTCTTATCTCCCTCTTCTGGCTCTCACCCCTCT -----+-----+-----+-----+-----+-----+-----+	720
721	L F L R K L H C T R N Y I H M N L F A S CTTGTTTCTTCGAAAACCTCCACTGCACGCGCAACTACATCCACATGAACCTGTTTGCTTC -----+-----+-----+-----+-----+-----+-----+	780
781	F I L R T L A V L V K D V V F Y N S Y S TTTCATCCTGAGAACCTGGCTGTACTGGTGAAGGACGTCGTCTTCTACAACTCTTACTC -----+-----+-----+-----+-----+-----+-----+	840
841	K R P D N E N G W M S Y L S E M S T S C CAAGAGGCCTGACAAATGAGAATGGGTGGATGTCCTACCTGTCAGAGATGTCCACCTCCTG -----+-----+-----+-----+-----+-----+-----+	900
901	R S V Q V L L H Y ? V G A N Y L W L L V CCGCTCAGTCCAGGTTCTCTTGCACTTCTTGTGGGTGCCAATTACTTATGGCTGCTGGT -----+-----+-----+-----+-----+-----+-----+	960
961	TGAAGGCCTCTACCTCCACACGCTGCTGGAGCCACAGTGCTTCCTGAGAGGCGGCTGTG -----+-----+-----+-----+-----+-----+-----+	1020
1021	P R Y L L L G W A F P V L F V V P W G F GCCCAGATACCTGCTGTTGGGTGGGCCCTTCCCTGTGCTATTGTTGTACCCTGGGGTTT -----+-----+-----+-----+-----+-----+-----+	1080
1080	A R A H L E N T G C W T T N G N K K I W CGCCCGTGCACACCTGGAGAACACAGGGTGCTGGACAACAAATGGGAATAAGAAAATCTG -----+-----+-----+-----+-----+-----+-----+	1140
1141	W I I R G P M M L C V T V N F F I F L K GTGGATCATCCGAGGACCCATGATGCTCTGTGTAACAGTCAATTTCTTCATCTTCTGAA -----+-----+-----+-----+-----+-----+-----+	1200
1201	I L K L L I S K L K A H Q M C F R D Y K AATTCTCAAGCTTCTCATTCTAAGCTCAAAGCTCATCAAATGTGCTTCAGAGATTATAA -----+-----+-----+-----+-----+-----+-----+	1250
1261	Y R L A K S T L V L I P L L G V H E I L ATACAGATTGGCAAAATCAACACTGGTCCTCATTCTTTATTGGGCGTTCATGAGATCCT -----+-----+-----+-----+-----+-----+-----+	1320
1321	F S F I T D D Q V E G F A K L I R L F I CTTCTCTTCATCACTGATGATCAAGTTGAAGGATTTGCAAACTTATACGACTTTTCAT -----+-----+-----+-----+-----+-----+-----+	1380

**FIG. 6B**

	Q L T L S S F H G F L V A L Q Y G F A N	
1381	TCAGTTGACACTGAGCTCCTTTTCATGGGTTCCTGGTGGCCTTGCAGTATGGTTTTGCCAA	1440
	G E V K A E L R K Y W V R F L L A R H S	
1441	TGGAGAAGTGAAGGCTGAGCTGCGGAAATACTGGGTCCGCTTCTTGCTAGCCCCGCACTC	1500
	G C R A C V L G K D F R F L G K C P K K	
1501	AGGCTGCAGAGCCTGTGTCTTGGGGAAGGACTTCCGGTTCCTAGGAAAATGTCCCAAGAA	1560
	L S E G D G A E K L R K L Q P S L N S G	
1561	GCTCTCGGAAGGAGATGGCGCTGAGAAGCTTCGGAAGCTGCAGCCCTCACTTAACAGTGG	1620
	R L L H L A M R G L G E L G A Q P Q Q D	
1621	GCGGCTCCTACATCTAGCCATGCGAGGTCTTGGGGAGCTGGGCGCCCAGCCCCAACAGGA	1680
	H A R W P R G S S L S E C S E G D V T M	
1681	CCATGCACGCTGGCCCCGGGGCAGCAGCCTGTCCGAGTGCAGTGAGGGGGATGTCACCAT	1740
	A N T M E E I L E E S E I	
1741	GGCCAACACCATGGAGGAGATTCTGGAAGAGAGTGAGATCTAGGGTGGAGTTCACCACC	1800
	CTGGCTCTGCTCCCAGGGACTCTTGAGGGGGCCCAGGAAGAGGAAGCAAGCAGGACACA	
1801		1860
	CGTTGCTGGGCACGGAATCATTCTCGTTCATTACCATGCCACTTTGATATGAAAGCTA	
1861		1920
	TCACAAGGTTCTTCAAGCTCTGTATGAAAGAGGCTGTGTGTCATGCTCACAGCCTCTGCC	
1921		1980
	TGCTCTTCTCATCCTAATAACCCCCACCAGTGTGTTTTCCACAATGCCACCAGACCCTA	
1981		2040
	GGGCCTGGCTCTAAATTCAAGCCAATGAAGTCCCAACCCGGAATTCTTTTGCTTTTACCC	
2041		2100
	CTGGAAGAAATA	
2101		2112

FIG. 6C

10/13

1 MKLGSSRAGP GRGSAGLLPG VHELPMGIPA PWGTSPLSFH RKCSLWAPGR  
51 PFLTLLVLLVS IKQVTGSLLE ETTRKWAQYK QACLRDLLKE PSGIFCNGTF  
101 DQYVCWPHSS PGNVSVPCPS YLPWWSEESS GRAYRHCLAQ GTWQTENAT  
151 DIWQDDSECS ENHSFKQNV DRYALLSTLQL MYTVGYFSLSL ISLFLALILL  
201 LFLRKLHCTR NYIHMLNFAS FILRTLAVLV KDVPFYNSYS KRPDNENGWM  
251 SYLSEMSTSC RSVQVLLHYF VGANYLWLLV EGYLHLLLE PTVLPERRLW  
301 PRYLLLGWAF PVLFVVPWGF ARAHLENTGC WTTNGNKKIW WIIRGPMMLC  
351 VTVNFFIFLK ILKLLISKLK AHQMCFRDYK YRLAKSTLVL IPLLGVHEIL  
401 FSFITDDQVE GFAKLIRLFI QLTLSFHFGL LVALQYGFAN GEVKAELRKY  
451 WVRFLARHS GCRACVLGKD FRFLGKCPKK LSEGDAEKL RKLQPSLNSG  
501 RLLHLAMRGL GELGAQPQD HARWPRGSSL SECSEGDVTM ANTMEEILEE  
551 SEI

FIG.7

11/13




FIG.8

12/13

1	MRPQSPAVPSRCREAPVPRVRAQPVGIPEAQGPVPLHSQMRLLWGPR	50
1	MKLGSSRAGPGRGSAGLLPGVHELPMGIPAPWGTSPLSFHRKCSLWAPGR	50
51	PFLALLLLVSIKQVTGSLLKETTKWANYKEKCLEDLHNRLSGIFCNGTF	100
51	PFLTLLVLLVSIKQVTGSLLLEETTRKWAQYKQACLRDLLKEPSGIFCNGTF	100
101	DRYVCWPHSYPGNVSVPCPSYLPWWNAESPGRAYRHCLAQGTWOTRENTT	150
101	DQYVCWPHSSPGNVSVPCPSYLPWWSEESSGRAYRHCLAQGTWOTIENAT	150
151	DIWQDESECSSENHSFRQNVHDYALLYTLQLMYTVGYSVSLISLFLALTFL	200
151	DIWQDDSECSSENHSFKQNVDRYALLSTLQLMYTVGYSFSLISLFLALTLL	200
201	LFLRKLHCTRNYIHMNLFASFILKVLAVLVKDMVSHNSYSKRPDDESGWM	250
201	LFLRKLHCTRNYIHMNLFASFILRTLAVLVKDVVFYNSYSKRPDNENGWM	250
251	SYLSETSVSCRSVQVLLHYFVGTHNLWLLVEGLYLHTLLEPTVFPERRLW	300
251	SYLSEMSTSCRSVQVLLHYFVGANYLWLLVEGLYLHTLLEPTVLPERRLW	300
301	PKYL VVGWAF PMLFVIPWGF ARAHLENTRCWATNGNLKIWWIIRGPMMLC	350
301	PRYLLLGWAF PVLFPVPWGF ARAHLENTGCWTNGNKKIWWIIRGPMMLC	350
351	VTVNFFIFLKILKLLISKLAHQMCFRDYKYRLAKSTLLLIPLLGVHEVL	400
351	VTVNFFIFLKILKLLISKLAHQMCFRDYKYRLAKSTLVLIPLLGVHEIL	400
401	FTFFPDDQVQGF SKRIRLF IQLTLSSVHGFLVALQYGFANGEVKAELRKS	450
401	FSFITDDQVEGFAKLIRLF IQLTLSSFHGFLVALQYGFANGEVKAELRKY	450
451	WGRFLLARHMGCRCTVLGKNFRFLGKCSKKLSEGDGSETLQKLRFSTCSS	500
451	WVRFLLARHSGCRACVLGKDFRFLGKCPKKLSEGDAEKLRLQPSLNSG	500
501	...HLASETLGDVGVPHRGRGANPRGSSLSSESSEGDFTLANTMEEILEE	547
501	RLLHLAMRGLGELGAQPQDHARWPRGSSLSECSEGDVTMANTMEEILEE	550
548	SEI 550	
551	SEI 553	

FIG.9



13/13

GL2R\_RAT\_R  
HWBRPAT\_TR  
GLPR\_HUMAN

MRPQSPAVPSRCREAPVPRVRAQPVGIPEAQGPVPLHSQQMRLWGPGRPFLLALLLV  
MKLGSSRAGPGRGSAGLLPGVHELPMGIAPAWGTSPLSFHRKCSLWAPGRPFLLVLLV  
-----MAGAPGLRLALLLGMVGRAGPRP-----

\* \* \*

GL2R\_RAT\_R  
HWBRPAT\_TR  
GLPR\_HUMAN

SIKQVTGSLLKETTQKWANYKEKCLEDLHNRL---SGIFCNGTFDRYVCWPHSYPG-NVS  
SIKQVTGSLLLEETTRKWAQYKQACLRDLLKEP---SGIFCNGTFDRYVCWPHSYPG-NVS  
---QGATVSLWETVQKWREYRQCQSLTEDPPATDLFCNRTFDEYACWPDGEGPSFVN

\* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \*

GL2R\_RAT\_R  
HWBRPAT\_TR  
GLPR\_HUMAN

VPCPSYLPWWNAESPGRAYRHCLAQGTWQTRENTTDIWQDESECSENHFRQNVHDYALL  
VPCPSYLPWWSESSGRAYRHCLAQGTWQTIENATDIWQDSECSENHFRQNVDRYALL  
VSCPWYLPWASSVPQGHVYRFCTAEGWLQKDNSSLPWRDLSECSESKRGERSSPEEQLL

\* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \*

GL2R\_RAT\_R  
HWBRPAT\_TR  
GLPR\_HUMAN

YTLQLMYTVGYSVSLISLFLALTFLFLRKLHCTRNYIHMNLFASFILKVLAVLVKDMVS  
STLQLMYTVGYFSLSLFLALTLLFLRKLHCTRNYIHMNLFASFILRTLAVLVKDVVF  
F-LYIITYVGYSLSALVIASAILLGFRHLHCTRNYIHLNLFASFILRLSVFIKDAAL

\* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \*

GL2R\_RAT\_R  
HWBRPAT\_TR  
GLPR\_HUMAN

HNSYSKRPDDESGWMSYLS-ETSVSCRSVQVLLHYFVGINHLWLLVEGLYLHTLLEPTVF  
YNSYSKRPDNENGWMSYLS-EMSTSCRSVQVLLHYFVGANYLWLLVEGLYLHTLLEPTVL  
KWMYST-AAQHQWDGLLSYQDSLSCRLVFLMQYCVAANYWLLVEGVLYTLAFSVF

\* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \*

GL2R\_RAT\_R  
HWBRPAT\_TR  
GLPR\_HUMAN

PERRLWPKYLVGWAFPMFLVIPWGFARAHLENTRCWATNGNLKIWWIIRGPMMLCVTN  
PERRLWFRYLLGWAFVFLVVPWGFARAHLENTGCWTNGNKKIWWIIRGPMMLCVTN  
SEQWIFRLYVSIGWGVPLLFVVPWGVKYLVEDEGCWTRNSNMNYWLIIRLPILFAIGVN

\* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \*

GL2R\_RAT\_R  
HWBRPAT\_TR  
GLPR\_HUMAN

FFIFLKILKLLISKLAHQMCFRDYKYRLAKSTLLLIPLLGVHEVLF TFFPDDQVQGF SK  
FFIFLKILKLLISKLAHQMCFRDYKYRLAKSTLVIPLLGVHEILFSITDDQVEGFAK  
FLIFVRVICIVSKLKANLMCKTDIKCRLAKSTLTLIPLLGTHEVIFAFVMDHARGTLR

\* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \*

GL2R\_RAT\_R  
HWBRPAT\_TR  
GLPR\_HUMAN

RIRLFIQLTLSSVHGF LVALQYGFANGEVKAELRKSWGRFLLARHWCRCV LGKNFRFL  
LIRLFIQLTLSSFHGF LVALQYGFANGEVKABLRKYKWRFLARHSGCRACV LGDRFRFL  
FIKLFTELSTSFQGLMVAI LYCFVNNEVQLEFRKSWERWRLHHLIQRDSSMKP-----

\* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \* \*

GL2R\_RAT\_R  
HWBRPAT\_TR  
GLPR\_HUMAN

GKCSKKLSEGDGSETLQKLRFTSCSS--HLASETLGDVGVPHRGRGAWPRGSSLSESS  
GKCPKKLSEGDGAELRKLPQSLNSGRLLHLMRGLGELGAQPQDQDARWPRGSSLSECS  
LKCP TSS-LSSGATAGSSMYTATCQASCS-----

\* \* \* \* \*

GL2R\_RAT\_R  
HWBRPAT\_TR  
GLPR\_HUMAN

EGDFTLANTMEEILEESEI  
EGDVTMANTMEEILEESEI  
-----

FIG.10

**CORRECTED  
VERSION\***

**PCT**

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : C12N 15/12, C07K 14/705, C12N 5/10, 15/62, C07K 16/28, C12Q 1/68, G01N 33/50		A3	(11) International Publication Number: <b>WO 98/25955</b> (43) International Publication Date: 18 June 1998 (18.06.98)
(21) International Application Number: PCT/CA97/00969 (22) International Filing Date: 15 December 1997 (15.12.97) (30) Priority Data: 08/767,224 13 December 1996 (13.12.96) US 08/787,721 24 January 1997 (24.01.97) US 08/845,546 24 April 1997 (24.04.97) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications US 08/767,224 (CIP) Filed on 13 December 1996 (13.12.96) US 08/787,721 (CIP) Filed on 24 January 1997 (24.01.97) US 08/845,546 (CIP) Filed on 24 April 1997 (24.04.97) (71) Applicant (for all designated States except US): ALLELIX BIOPHARMACEUTICALS INC. [CA/CA]; 6850 Goreway Drive, Mississauga, Ontario L4V 1V7 (CA).		(72) Inventors; and (75) Inventors/Applicants (for US only): MUNROE, Donald, G. [CA/CA]; 27 Wakefield Lane, Waterdown, Ontario L0R 2H3 (CA). GUPTA, Ashwani, K. [CA/CA]; 7031 Dunrobin Way, Mississauga, Ontario L5N 6Y4 (CA). VYAS, Tejal, B. [CA/CA]; 275 Riel Drive, Mississauga, Ontario L5B 3K1 (CA). McCALLUM, Kirk [CA/CA]; 3918 Pitch Pine Crescent, Mississauga, Ontario L5L 1P8 (CA). FAN, Ermei [CA/CA]; Apartment #418, 20 Southport Street, Toronto, Ontario M6S 4Y8 (CA). (74) Agent: AITKEN, David, W.; Osler, Hoskin & Harcourt, Suite 1500, 50 O'Connor Street, Ottawa, Ontario K1P 6L2 (CA). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
		<b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limits for amending the claims</i> <i>and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 30 July 1998 (30.07.98)	

(54) Title: CLONED GLUCAGON-LIKE PEPTIDE-2 RECEPTORS

(57) Abstract

The invention relates to nucleotides and amino acid sequences encoding glucagon-like peptide 2 receptors, recombinant host cells transformed with such nucleotides, and methods of using the same in drug screening and related applications.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/CA 97/00969

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C07K14/705 C12N5/10 C12N15/62 C07K16/28  
C12Q1/68 G01N33/50

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K C12N C12Q G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DRUCKER D J ET AL: "INDUCTION OF INTESTINAL EPITHELIAL PROLIFERATION BY GLUCAGON-LIKE PEPTIDE 2" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 93, 1990, pages 7911-7916, XP000650124	19
Y	whole document, especially page 7914, right column + page 7915	1-5, 12, 13
X	WO 96 32414 A (1149336 ONTARIO INC) 17 October 1996 pages 2, 3, 6, 7	19
Y	WO 93 19175 A (THORENS BERNARD) 30 September 1993 abstract, examples 1-3	1-5, 12, 13
	--- -/-	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

4 June 1998

Date of mailing of the international search report

10/06/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Holtorf, S

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 97/00969

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO 95 04821 A (MERCK &amp; CO INC ;GRAZIANO  MICHAEL P (US); BORKOWSKI DOREEN A (US);)  16 February 1995  pages 1-3, page 5 , line 27-33; page 6;  example 5</p> <p>-----</p>	1-19

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 97/00969

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9632414 A	17-10-1996	AU 5265896 A EP 0830377 A	30-10-1996 25-03-1998
WO 9319175 A	30-09-1993	AU 3751293 A DE 586657 T EP 0586657 A ES 2070801 T JP 6508765 T US 5670360 A	21-10-1993 18-01-1996 16-03-1994 16-06-1995 06-10-1994 23-09-1997
WO 9504821 A	16-02-1995	EP 0716694 A JP 9501054 T	19-06-1996 04-02-1997